

# thm\_2Einteger\_2EINT\_\_NOT\_\_LE (TMFid4giVtdnJPaTkEG1ejTii3jMQNVTAkb)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Eenum\_2Eenum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eenum\_2Eenum \tag{1}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{2}$$

Let  $ty\_2Einteger\_2Eint : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Einteger\_2Eint \tag{3}$$

Let  $c\_2Einteger\_2Eint\_REP\_CLASS : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Eint\_REP\_CLASS \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Eenum\_2Eenum\ ty\_2Eenum\_2Eenum)})\ ty\_2Einteger\_2Eint) \tag{4}$$

**Definition 7** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define  $c\_2Einteger\_2Eint\_REP$  to be  $\lambda V0a \in ty\_2Einteger\_2Eint.(ap (c\_2Emin\_2E\_40 (ty\_2Einteger\_2Eint\_REP\_CLASS a)))$

Let  $c\_2Einteger\_2Etint\_lt : \iota$  be given. Assume the following.

$$c\_2Einteger\_2Etint\_lt \in ((2^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)})^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum)}) \quad (5)$$

**Definition 9** We define  $c\_2Einteger\_2Eint\_lt$  to be  $\lambda V0T1 \in ty\_2Einteger\_2Eint.\lambda V1T2 \in ty\_2Einteger\_2Eint.$

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_7E))$

**Definition 11** We define  $c\_2Einteger\_2Eint\_le$  to be  $\lambda V0x \in ty\_2Einteger\_2Eint.\lambda V1y \in ty\_2Einteger\_2Eint.$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

**Theorem 1**

$$(\forall V0x \in ty\_2Einteger\_2Eint.(\forall V1y \in ty\_2Einteger\_2Eint.((\neg(p\ (ap\ (ap\ c\_2Einteger\_2Eint\_le\ V0x)\ V1y))) \Leftrightarrow (p\ (ap\ (ap\ c\_2Einteger\_2Eint\_lt\ V1y)\ V0x))))))$$