

thm_2Einteger_2EINT__POS__NZ (TMMvoBzg- BfLMV4YNmMdFiVatKUKjtTPXcAs)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{4}$$

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint^{ty_2Enum_2Enum}) \tag{5}$$

Definition 3 We define $c_2Ebool_2E_2ET$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_2E21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1Q \in 2.V1Q))\ (\lambda V2R \in 2.V2R))\ P)\ a$

Definition 5 We define $c_2Ebool_2E_2EF$ to be $(ap\ (c_2Ebool_2E_2E21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 6 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 7 We define $c_2Ebool_2E_2E7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_2EF))$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (6)$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \quad (7)$$

Definition 8 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 9 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E40\ (ty_2Einteger_2Eint_REP_CLASS\ V0a)))$

Let $c_2Einteger_2Eint_lt : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (8)$$

Definition 10 We define $c_2Einteger_2Eint_lt$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.\lambda V1T2 \in ty_2Einteger_2Eint.$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a.(\forall V1y \in A.27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (9)$$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.(\forall V1y \in ty_2Einteger_2Eint.((p\ (ap\ (ap\ c_2Einteger_2Eint_lt\ V0x)\ V1y)) \Rightarrow (\neg(V0x = V1y)))))) \quad (10)$$

Theorem 1

$$(\forall V0x \in ty_2Einteger_2Eint.((p\ (ap\ (ap\ c_2Einteger_2Eint_lt\ (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0))\ V0x)) \Rightarrow (\neg(V0x = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0))))))$$