

thm_2Einteger_2EINT__QUOT__CALCULATE
(TMT-
FimShJ6uzX3meCn94F7CTpdfQm93z1wL)

October 26, 2020

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Earithmetic_2EDIV : \iota$ be given. Assume the following.

$$c_2Earithmetic_2EDIV \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{2}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{3}$$

Let $ty_2Einteger_2Eint : \iota$ be given. Assume the following.

$$nonempty\ ty_2Einteger_2Eint \tag{4}$$

Let $c_2Einteger_2Eint_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{ty_2Einteger_2Eint}) \tag{5}$$

Definition 1 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$).

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_21$ to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_27a$ to be $\lambda A.\lambda P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a})))$

Definition 5 We define $c_2Einteger_2Eint_REP$ to be $\lambda V0a \in ty_2Einteger_2Eint.(ap\ (c_2Emin_2E_40\ (ty_2Einteger_2Eint\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a})))$

Let $c_2Einteger_2Etint_neg : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_neg \in ((ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum) (ty_2Enum_2Enum))^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)} \quad (6)$$

Let $c_2Einteger_2Etint_eq : \iota$ be given. Assume the following.

$$c_2Einteger_2Etint_eq \in ((2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum)}) \quad (7)$$

Let $c_2Einteger_2Eint_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_ABS_CLASS \in (ty_2Einteger_2Eint)^{(2^{(ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)})} \quad (8)$$

Definition 6 We define $c_2Einteger_2Eint_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod\ ty_2Enum_2Enum\ ty_2Enum_2Enum)$

Definition 7 We define $c_2Einteger_2Eint_neg$ to be $\lambda V0T1 \in ty_2Einteger_2Eint.(ap\ c_2Einteger_2Eint_neg\ T1)$

Let $c_2Einteger_2Eint_quot : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_quot \in ((ty_2Einteger_2Eint)^{ty_2Einteger_2Eint})^{ty_2Einteger_2Eint} \quad (9)$$

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ t2)\ t1))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (10)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum)^{\omega} \quad (11)$$

Definition 10 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Einteger_2Eint_of_num : \iota$ be given. Assume the following.

$$c_2Einteger_2Eint_of_num \in (ty_2Einteger_2Eint)^{ty_2Enum_2Enum} \quad (12)$$

Definition 11 We define $c_2Ebool_2E_2F$ to be $(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V0t \in 2.V0t))$.

Definition 12 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap\ (ap\ c_2Emin_2E_3D_3D_3E\ V0t)\ c_2Ebool_2E_7E))$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.((ap\ c_2Einteger_2Eint_neg\ (ap\ c_2Einteger_2Eint_neg\ V0x)) = V0x)) \quad (13)$$

Assume the following.

$$(\forall V0x \in ty_2Einteger_2Eint.(((ap\ c_2Einteger_2Eint_neg\ V0x) = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) \Leftrightarrow (V0x = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)))) \quad (14)$$

Assume the following.

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.((ap\ c_2Einteger_2Eint_of_num\ V0m) = (ap\ c_2Einteger_2Eint_of_num\ V1n)) \Leftrightarrow (V0m = V1n)))) \quad (15)$$

Assume the following.

$$(\forall V0p \in ty_2Enum_2Enum.(\forall V1q \in ty_2Enum_2Enum.(\neg(V1q = c_2Enum_2E0)) \Rightarrow ((ap\ (ap\ c_2Einteger_2Eint_quot\ (ap\ c_2Einteger_2Eint_of_num\ V0p))\ (ap\ c_2Einteger_2Eint_of_num\ V1q)) = (ap\ c_2Einteger_2Eint_of_num\ (ap\ (ap\ c_2Arithmetic_2EDIV\ V0p)\ V1q)))))) \quad (16)$$

Assume the following.

$$(\forall V0p \in ty_2Einteger_2Eint.(\forall V1q \in ty_2Einteger_2Eint.(\neg(V1q = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) \Rightarrow (((ap\ (ap\ c_2Einteger_2Eint_quot\ (ap\ c_2Einteger_2Eint_neg\ V0p))\ V1q) = (ap\ c_2Einteger_2Eint_neg\ (ap\ (ap\ c_2Einteger_2Eint_quot\ V0p)\ V1q))) \wedge ((ap\ (ap\ c_2Einteger_2Eint_quot\ V0p)\ (ap\ c_2Einteger_2Eint_neg\ V1q)) = (ap\ c_2Einteger_2Eint_neg\ (ap\ (ap\ c_2Einteger_2Eint_quot\ V0p)\ V1q))))))) \quad (17)$$

Theorem 1

$$((\forall V0p \in ty_2Enum_2Enum.(\forall V1q \in ty_2Enum_2Enum.(\neg(V1q = c_2Enum_2E0)) \Rightarrow ((ap\ (ap\ c_2Einteger_2Eint_quot\ (ap\ c_2Einteger_2Eint_of_num\ V0p))\ (ap\ c_2Einteger_2Eint_of_num\ V1q)) = (ap\ c_2Einteger_2Eint_of_num\ (ap\ (ap\ c_2Arithmetic_2EDIV\ V0p)\ V1q)))))) \wedge ((\forall V2p \in ty_2Einteger_2Eint.(\forall V3q \in ty_2Einteger_2Eint.(\neg(V3q = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) \Rightarrow (((ap\ (ap\ c_2Einteger_2Eint_quot\ (ap\ c_2Einteger_2Eint_neg\ V2p))\ V3q) = (ap\ c_2Einteger_2Eint_neg\ (ap\ (ap\ c_2Einteger_2Eint_quot\ V2p)\ V3q))) \wedge ((ap\ (ap\ c_2Einteger_2Eint_quot\ V2p)\ (ap\ c_2Einteger_2Eint_neg\ V3q)) = (ap\ c_2Einteger_2Eint_neg\ (ap\ (ap\ c_2Einteger_2Eint_quot\ V2p)\ V3q))))))) \wedge ((\forall V4m \in ty_2Enum_2Enum.(\forall V5n \in ty_2Enum_2Enum.((ap\ c_2Einteger_2Eint_of_num\ V4m) = (ap\ c_2Einteger_2Eint_of_num\ V5n)) \Leftrightarrow (V4m = V5n)))) \wedge ((\forall V6x \in ty_2Einteger_2Eint.(((ap\ c_2Einteger_2Eint_neg\ V6x) = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)) \Leftrightarrow (V6x = (ap\ c_2Einteger_2Eint_of_num\ c_2Enum_2E0)))) \wedge ((\forall V7x \in ty_2Einteger_2Eint.((ap\ c_2Einteger_2Eint_neg\ (ap\ c_2Einteger_2Eint_neg\ V7x)) = V7x))))))$$