

thm_2Eintegral_2EINTEGRABLE__CMUL
(TMN4RJsXb5rt6SLDEbNfBtvHsf9LLQ8av4C)

October 26, 2020

Let $ty_2Ehreal_2Ehreal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Ehreal_2Ehreal \tag{1}$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \tag{2}$$

Let $ty_2Erealax_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealax_2Ereal \tag{3}$$

Let $c_2Erealax_2Ereal_REP_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_REP_CLASS \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ ty_2Erealax_2Ereal) \tag{4}$$

Definition 1 We define c_2Emin_2E40 to be $\lambda A.\lambda P \in 2^A$.if $(\exists x \in A.p\ (ap\ P\ x))$ **then** (the $(\lambda x.x \in A \wedge p\ x)$ of type $\iota \Rightarrow \iota$).

Definition 2 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o\ (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define c_2Ebool_2E27 to be $(ap\ (ap\ (c_2Emin_2E3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define c_2Ebool_2E21 to be $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x))\ P))$

Definition 5 We define $c_2Erealax_2Ereal_REP$ to be $\lambda V0a \in ty_2Erealax_2Ereal.(ap\ (c_2Emin_2E40\ (ty_2Erealax_2Ereal_REP_CLASS\ a)))$

Let $c_2Erealax_2Etrealt_lt : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealt_lt \in ((2^{(ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal\ ty_2Ehreal_2Ehreal)})\ (ty_2Epair_2Eprod\ ty_2Ehreal_2Ehreal)) \tag{5}$$

Definition 6 We define $c_2Erealax_2Ereal_lt$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$

Definition 7 We define $c_Ebool_2E_2F$ to be $(ap (c_Ebool_2E_21) 2) (\lambda V0t \in 2.V0t)$.

Definition 8 We define $c_Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define $c_Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_Emin_2E_3D_3D_3E V0t) c_Ebool_2E_2F))$.

Definition 10 We define $c_Ereal_2Ereal_lte$ to be $\lambda V0x \in ty_2Erealax_2Ereal.\lambda V1y \in ty_2Erealax_2Ereal$.

Definition 11 We define $c_Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_Ebool_2E_21) 2) (\lambda V2t \in 2.V2t)))$.

Definition 12 We define $c_Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_Emin_2E_40$

Let $c_2Eintegral_2Eintegral : \iota$ be given. Assume the following.

$$c_2Eintegral_2Eintegral \in ((ty_2Erealax_2Ereal^{(ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}})})^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Ereal)}})) \quad (6)$$

Let $c_2Eintegral_2Eintegrable : \iota$ be given. Assume the following.

$$c_2Eintegral_2Eintegrable \in ((2^{(ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal})^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Ereal)}}))^{(ty_2Epair_2Eprod ty_2Erealax_2Ereal ty_2Ereal)}})) \quad (7)$$

Let $c_2Erealax_2Etrealmul : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealmul \in (((ty_2Epair_2Eprod ty_2Ehreal_2Ehreal^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}}))^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}}))^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}})) \quad (8)$$

Let $c_2Erealax_2Etrealeq : \iota$ be given. Assume the following.

$$c_2Erealax_2Etrealeq \in ((2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}}))^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}})) \quad (9)$$

Let $c_2Erealax_2Ereal_ABS_CLASS : \iota$ be given. Assume the following.

$$c_2Erealax_2Ereal_ABS_CLASS \in (ty_2Erealax_2Ereal^{(2^{(ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)}})}) \quad (10)$$

Definition 13 We define $c_2Erealax_2Ereal_ABS$ to be $\lambda V0r \in (ty_2Epair_2Eprod ty_2Ehreal_2Ehreal ty_2Ehreal_2Ehreal)$.

Definition 14 We define $c_2Erealax_2Ereal_mul$ to be $\lambda V0T1 \in ty_2Erealax_2Ereal.\lambda V1T2 \in ty_2Erealax_2Ereal$.

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2Epair_2EABS_prod A_27a A_27b \in ((ty_2Epair_2Eprod A_27a A_27b)^{(2^{A_27b})^{A_27a}}) \quad (11)$$

Definition 15 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0x \in A_27a.\lambda V1y \in A_27b.(ap (c_2E$

Let $c_2Etransc_2EDint : \iota$ be given. Assume the following.

$$c_2Etransc_2EDint \in (((2^{ty_2Erealax_2Ereal})(ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal})))(ty_2Epair_2Eprod\ ty_2Erealax_2Ereal) \quad (12)$$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (14)$$

Assume the following.

$$(\forall V0a \in ty_2Erealax_2Ereal. (\forall V1b \in ty_2Erealax_2Ereal. (\forall V2f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}). ((p\ (ap\ (ap\ c_2Eintegral_2Eintegrable\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V0a)\ V1b))\ V2f)) \Leftrightarrow (\exists V3i \in ty_2Erealax_2Ereal. (p\ (ap\ (ap\ (ap\ c_2Etransc_2EDint\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V0a)\ V1b))\ V2f)\ V3i)))))) \quad (15)$$

Assume the following.

$$(\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}). (\forall V1a \in ty_2Erealax_2Ereal. (\forall V2b \in ty_2Erealax_2Ereal. ((p\ (ap\ (ap\ c_2Eintegral_2Eintegrable\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V1a)\ V2b))\ V0f)) \Rightarrow (p\ (ap\ (ap\ (ap\ c_2Etransc_2EDint\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V1a)\ V2b))\ V0f)\ (ap\ (ap\ c_2Eintegral_2Eintegral\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V1a)\ V2b))\ V0f)))))) \quad (16)$$

Assume the following.

$$(\forall V0f \in (ty_2Erealax_2Ereal^{ty_2Erealax_2Ereal}). (\forall V1a \in ty_2Erealax_2Ereal. (\forall V2b \in ty_2Erealax_2Ereal. (\forall V3c \in ty_2Erealax_2Ereal. (\forall V4i \in ty_2Erealax_2Ereal. ((p\ (ap\ (ap\ c_2Etransc_2EDint\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V1a)\ V2b))\ V0f)\ V4i)) \Rightarrow (p\ (ap\ (ap\ (ap\ c_2Etransc_2EDint\ (ap\ (ap\ (c_2Epair_2E_2C\ ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal)\ V1a)\ V2b))\ (\lambda V5x \in ty_2Erealax_2Ereal. (ap\ (ap\ c_2Erealax_2Ereal_mul\ V3c)\ (ap\ V0f\ V5x))))\ (ap\ (ap\ c_2Erealax_2Ereal_mul\ V3c)\ V4i)))))) \quad (17)$$

Theorem 1

$$\begin{aligned} & (\forall V0f \in (ty_2Erealax_2Ereal\ ty_2Erealax_2Ereal).(\forall V1a \in \\ & ty_2Erealax_2Ereal.(\forall V2b \in ty_2Erealax_2Ereal.(\forall V3c \in \\ & ty_2Erealax_2Ereal.(((p (ap (ap (c_2Ereal_2Ereal_lte V1a) V2b)) \wedge \\ (p (ap (ap (c_2Eintegral_2Eintegrable (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal \\ ty_2Erealax_2Ereal) V1a) V2b)) V0f)))) \Rightarrow (p (ap (ap (c_2Eintegral_2Eintegrable \\ (ap (ap (c_2Epair_2E_2C ty_2Erealax_2Ereal ty_2Erealax_2Ereal) \\ V1a) V2b)) (\lambda V4x \in ty_2Erealax_2Ereal.(ap (ap (c_2Erealax_2Ereal_mul \\ V3c) (ap V0f V4x)))))))))))))) \end{aligned}$$