

thm\_2Eintegral\_2ELE\_0  
 (TMGT3uErruuMmdpW45gpeD8WQiVnXCqe6jD)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (1)$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A. \lambda x \in A. \lambda y \in A. inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P)))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2. \lambda Q \in 2. inj\_o (p \ P \Rightarrow p \ Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF))$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2. inj\_o (V0t1 = V1t2))))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \quad (2)$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \quad (3)$$

Let  $c\_2Enum\_2EAABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EAABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \quad (4)$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. (ap c\_2Enum\_2EAABS\_num m)$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum. \lambda V1n \in ty\_2Enum\_2Enum.$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (5)$$

**Definition 14** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP).$

Assume the following.

$$\begin{aligned} ((\forall V0n \in ty\_2Enum\_2Enum.((p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D \\ V0n)\ c\_2Enum\_2E0)) \Leftrightarrow (V0n = c\_2Enum\_2E0))) \wedge (\forall V1m \in ty\_2Enum\_2Enum. \\ (\forall V2n \in ty\_2Enum\_2Enum.((p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D \\ V1m)\ (ap\ c\_2Enum\_2ESUC\ V2n))) \Leftrightarrow ((V1m = (ap\ c\_2Enum\_2ESUC\ V2n)) \vee \\ (p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V1m)\ V2n))))))) \end{aligned} \quad (6)$$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \vee (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \vee True) \Leftrightarrow True) \wedge \\ (((False \vee (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee False) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \vee \\ (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall A\_27a.\ nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (9)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} (\forall V0P \in (2^{ty\_2Enum\_2Enum}).(((p\ (ap\ V0P\ c\_2Enum\_2E0)) \wedge \\ (\forall V1n \in ty\_2Enum\_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c\_2Enum\_2ESUC\ \\ V1n)))))) \Rightarrow (\forall V2n \in ty\_2Enum\_2Enum.(p\ (ap\ V0P\ V2n)))))) \end{aligned} \quad (11)$$

### Theorem 1

$$(\forall V0n \in ty\_2Enum\_2Enum.(p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D \\ c\_2Enum\_2E0)\ V0n)))$$