

thm\_2Eintegral\_2ESUM\_EQ\_0  
(TMafuchyunJu6uyzTnqyYPbNBpPqx1bAA9J)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F))$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{2}$$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (omega^{ty\_2Enum\_2Enum}) \tag{3}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (omega^{omega}) \tag{4}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{omega}) \tag{5}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p\ (ap\ P\ x)) \text{ then } (the\ (\lambda x.x \in A \wedge p$   
of type  $\iota \Rightarrow \iota$ .

**Definition 10** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 13** We define  $c\_2Earithmetic\_2E\_3C\_3D$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{6}$$

**Definition 14** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $ty\_2Erealx\_2Ereal : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Erealx\_2Ereal \tag{7}$$

Let  $c\_2Ereal\_2Ereal\_of\_num : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Ereal\_of\_num \in (ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum}) \tag{8}$$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Epair\_2Eprod\ A0\ A1) \tag{9}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epair\_2EABS\_prod\ A\_27a\ A\_27b \in ((ty\_2Epair\_2Eprod\ A\_27a\ A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{10}$$

**Definition 15** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap\ (c\_2$

Let  $c\_2Ereal\_2Esum : \iota$  be given. Assume the following.

$$c\_2Ereal\_2Esum \in ((ty\_2Erealx\_2Ereal^{(ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum})})^{(ty\_2Epair\_2Eprod\ ty\_2Enum\_2Enum)}) \tag{11}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.( \\ & (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V0m)\ V1n) = (ap\ (ap\ c\_2Earithmetic\_2E\_2B \\ & \quad V1n)\ V0m)))) \end{aligned} \tag{12}$$

Assume the following.

$$True \quad (13)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ \forall V0t1 \in A\_27a. (\forall V1t2 \in A\_27b. ((ap\ (\lambda V2x \in A\_27b. \\ V0t1)\ V1t2) = V0t1))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in \\ A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow \\ True)) \end{aligned} \quad (16)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in \\ A\_27a. (\forall V2z \in A\_27a. (((V0x = V1y) \wedge (V1y = V2z)) \Rightarrow (V0x = V2z)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg( \\ p\ V0t)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} (\forall V0f \in (ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum}). (\forall V1g \in \\ (ty\_2Erealx\_2Ereal^{ty\_2Enum\_2Enum}). (\forall V2m \in ty\_2Enum\_2Enum. \\ (\forall V3n \in ty\_2Enum\_2Enum. ((\forall V4r \in ty\_2Enum\_2Enum. \\ (((p\ (ap\ (ap\ c\_2Earithmetic\_2E\_3C\_3D\ V2m)\ V4r)) \wedge (p\ (ap\ (ap\ c\_2Eprim\_rec\_2E\_3C \\ V4r)\ (ap\ (ap\ c\_2Earithmetic\_2E\_2B\ V3n)\ V2m)))))) \Rightarrow ((ap\ V0f\ V4r) = ( \\ ap\ V1g\ V4r)))))) \Rightarrow ((ap\ (ap\ c\_2Ereal\_2Esum\ (ap\ (ap\ (c\_2Epair\_2E\_2C \\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)\ V2m)\ V3n))\ V0f) = (ap\ (ap\ c\_2Ereal\_2Esum \\ (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Enum\_2Enum\ ty\_2Enum\_2Enum)\ V2m)\ V3n)) \\ V1g)))))) \end{aligned} \quad (19)$$

Assume the following.

$$\begin{aligned} (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ (ap\ (ap\ c\_2Ereal\_2Esum\ (ap\ (ap\ (c\_2Epair\_2E\_2C\ ty\_2Enum\_2Enum \\ ty\_2Enum\_2Enum)\ V0m)\ V1n))\ (\lambda V2r \in ty\_2Enum\_2Enum. (ap\ c\_2Ereal\_2Ereal\_of\_num \\ c\_2Enum\_2E0))) = (ap\ c\_2Ereal\_2Ereal\_of\_num\ c\_2Enum\_2E0)))) \end{aligned} \quad (20)$$

**Theorem 1**

$$\begin{aligned} & (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\ & \quad \forall V2f \in (ty\_2Erealax\_2Ereal^{ty\_2Enum\_2Enum}). ((\forall V3r \in \\ & \quad ty\_2Enum\_2Enum. (((p (ap (ap c\_2Earithmetic\_2E\_3C\_3D V0m) V3r)) \wedge \\ & \quad (p (ap (ap c\_2Eprim\_rec\_2E\_3C V3r) (ap (ap c\_2Earithmetic\_2E\_2B \\ & V0m) V1n)))) \Rightarrow ((ap V2f V3r) = (ap c\_2Ereal\_2Ereal\_of\_num c\_2Enum\_2E0)))) \Rightarrow \\ & \quad ((ap (ap c\_2Ereal\_2Esum (ap (ap (c\_2Epair\_2E\_2C ty\_2Enum\_2Enum \\ & ty\_2Enum\_2Enum) V0m) V1n)) V2f) = (ap c\_2Ereal\_2Ereal\_of\_num \\ & \quad c\_2Enum\_2E0)))))) \end{aligned}$$