

# thm\_2Eiterate\_2ENSUM\_DELETE (TMX- HTfp2vwrgFMAfKSkdgWS1qKmmbzfpMUo)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_2E1$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2E7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2EF))$

**Definition 7** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2E\_2EF)$ .

**Definition 8** We define  $c\_2Ebool\_2E\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 9** We define  $c\_2Ebool\_2E\_2E5C\_2EF$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V2t \in 2.V2t)))$

**Definition 10** We define  $c\_2Ebool\_2E\_2E2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \tag{1}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{2}$$

**Definition 11** We define  $c\_2Epair\_2E\_2E2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2Ebool\_2E\_2EIN$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC \\ A\_27a\ A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod\ A\_27a\ 2)^{A\_27b}}) \end{aligned} \quad (3)$$

**Definition 12** We define  $c\_2Epred\_set\_2EINSERT$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.\lambda V1s \in (2^{A\_27a}).(ap\ (c\_2E$

**Definition 13** We define  $c\_2Epred\_set\_2EDIFF$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap\ (c\_2E$

**Definition 14** We define  $c\_2Epred\_set\_2EDELETE$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1x \in A\_27a.(ap\ (ap$

**Definition 15** We define  $c\_2Epred\_set\_2EFINITE$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).(ap\ (c\_2Ebool\_2E\_21\ (2$

**Definition 16** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if}\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge P\ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 17** We define  $c\_2Eiterate\_2Eneutral$  to be  $\lambda A\_27a : \iota.\lambda V0op \in ((A\_27a^{A\_27a})^{A\_27a}).(ap\ (c\_2Emin$

**Definition 18** We define  $c\_2Eiterate\_2Esupport$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}).\lambda V$

**Definition 19** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.($

**Definition 20** We define  $c\_2Eiterate\_2EITSET$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in ((A\_27a^{A\_27a})^{A\_27b}).\lambda V$

**Definition 21** We define  $c\_2Eiterate\_2Eiterate$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}).\lambda V$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \quad (4)$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (5)$$

**Definition 22** We define  $c\_2Eiterate\_2Esum$  to be  $\lambda A\_27a : \iota.(ap\ (c\_2Eiterate\_2Eiterate\ A\_27a\ ty\_2Enum\_2Enum))$

**Definition 23** We define  $c\_2Eiterate\_2Emonoidal$  to be  $\lambda A\_27a : \iota.\lambda V0op \in ((A\_27a^{A\_27a})^{A\_27a}).(ap\ (ap\ c\_2E$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in \\ A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge \\
& (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t))))))
\end{aligned} \tag{8}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (( \\
& (p \ V0t) \Rightarrow False) \Leftrightarrow (\neg (p \ V0t))))))
\end{aligned} \tag{9}$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{10}$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{11}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\
& (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg (p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow (\neg ( \\
& p \ V0t))))))
\end{aligned} \tag{12}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p \ V0t1) \Rightarrow \\
& ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3))))))
\end{aligned} \tag{13}$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in \\
& 2.(((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))))) \Rightarrow \\
& (((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27))))))
\end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty \ A\_27a \Rightarrow \forall A\_27b.nonempty \ A\_27b \Rightarrow ( \\
& \forall V0op \in ((A\_27b^{A\_27b})^{A\_27b}).((p \ (ap \ (c\_2Eiterate\_2Emonoidal \\
& A\_27b) \ V0op)) \Rightarrow (\forall V1f \in (A\_27b^{A\_27a}).(\forall V2s \in (2^{A\_27a}). \\
& (\forall V3a \in A\_27a.(((p \ (ap \ (c\_2Epred\_set\_2EFINITE \ A\_27a) \ V2s)) \wedge \\
& (p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27a) \ V3a) \ V2s))) \Rightarrow ((ap \ (ap \ V0op \ (ap \ V1f \\
& V3a)) \ (ap \ (ap \ (ap \ (c\_2Eiterate\_2Eiterate \ A\_27a \ A\_27b) \ V0op) \ (ap \\
& (ap \ (c\_2Epred\_set\_2EDELETE \ A\_27a) \ V2s) \ V3a)) \ V1f)) = (ap \ (ap \ (ap \\
& (c\_2Eiterate\_2Eiterate \ A\_27a \ A\_27b) \ V0op) \ V2s) \ V1f))))))
\end{aligned} \tag{15}$$

Assume the following.

$$(p \ (ap \ (c\_2Eiterate\_2Emonoidal \ ty\_2Enum\_2Enum) \ c\_2Earithmetic\_2E\_2B)) \tag{16}$$

**Theorem 1**

$$\begin{aligned} & \forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0f \in (ty\_2Enum\_2Enum^{A_{.27a}}). \\ & (\forall V1s \in (2^{A_{.27a}}). (\forall V2a \in A_{.27a}. ((p (ap (c\_2Epred\_set\_2EFINITE \\ & A_{.27a}) V1s)) \wedge (p (ap (ap (c\_2Ebool\_2EIN A_{.27a}) V2a) V1s))) \Rightarrow ((ap \\ & (ap c\_2Earithmetic\_2E\_2B (ap V0f V2a)) (ap (ap (c\_2Eiterate\_2Ensum \\ & A_{.27a}) (ap (ap (c\_2Epred\_set\_2EDELETE A_{.27a}) V1s) V2a)) V0f)) = \\ & (ap (ap (c\_2Eiterate\_2Ensum A_{.27a}) V1s) V0f)))))) \end{aligned}$$