

# thm\_2Elist\_2ELIST\_\_BIND\_\_ID (TMVxRtc- npgmnhqnfKpG5Acm1aNWh24QwLYb)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Ecombin\_2EK$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.(\lambda V0x \in A\_27a.(\lambda V1y \in A\_27b.V0x))$

**Definition 6** We define  $c\_2Ecombin\_2ES$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda A\_27c : \iota.(\lambda V0f \in ((A\_27c^{A\_27b})^{A\_27a}))$

**Definition 7** We define  $c\_2Ecombin\_2EI$  to be  $\lambda A\_27a : \iota.(ap (ap (c\_2Ecombin\_2ES A\_27a (A\_27a^{A\_27a})) A\_27a))$

**Definition 8** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $ty\_2Elist\_2Elist : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Elist\_2Elist A0) \quad (1)$$

Let  $c\_2Elist\_2EMAP : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Elist\_2EMAP A\_27a A\_27b \in (((ty\_2Elist\_2Elist A\_27b)^{(ty\_2Elist\_2Elist A\_27a)})^{(A\_27b^{A\_27a})}) \quad (2)$$

Let  $c\_2Elist\_2EFLAT : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow c\_2Elist\_2EFLAT A\_27a \in ((ty\_2Elist\_2Elist A\_27a)^{(ty\_2Elist\_2Elist (ty\_2Elist\_2Elist A\_27a))}) \quad (3)$$

**Definition 10** We define  $c\_2Elist\_2ELIST\_BIND$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0l \in (ty\_2Elist\_2Elist A\_27a$

Assume the following.

$$True \tag{4}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \tag{5}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{6}$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0l \in (ty\_2Elist\_2Elist \\ & A\_27a).(((ap (ap (c\_2Elist\_2EMAP A\_27a A\_27a) (\lambda V1x \in A\_27a. \\ & V1x)) V0l) = V0l) \wedge ((ap (ap (c\_2Elist\_2EMAP A\_27a A\_27a) (c\_2Ecombin\_2El \\ & A\_27a)) V0l) = V0l))) \end{aligned} \tag{7}$$

**Theorem 1**

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0l \in (ty\_2Elist\_2Elist \\ & (ty\_2Elist\_2Elist A\_27a).(((ap (ap (c\_2Elist\_2ELIST\_BIND \\ & A\_27a (ty\_2Elist\_2Elist A\_27a)) V0l) (\lambda V1x \in (ty\_2Elist\_2Elist \\ & A\_27a).V1x)) = (ap (c\_2Elist\_2EFLAT A\_27a) V0l)) \wedge ((ap (ap (c\_2Elist\_2ELIST\_BIND \\ & A\_27a (ty\_2Elist\_2Elist A\_27a)) V0l) (c\_2Ecombin\_2El (ty\_2Elist\_2Elist \\ & A\_27a))) = (ap (c\_2Elist\_2EFLAT A\_27a) V0l)))) \end{aligned}$$