

thm_2Ellist_2ELFINITE_rules
(TMZGLmkt3qbg2AooSBhTELxSTQ1NUzgALxa)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 7 We define c_2Enum_2E0 to be $(ap c_2Enum_2EABS_num c_2Enum_2EZERO_REP)$.

Definition 8 We define $c_2Earithmic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 9 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 10 We define $c_2Earithmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic$

Definition 11 We define $c_2Earithmetic_2ENUMERAL$ to be $\lambda V0x \in ty_2Enum_2Enum.V0x$.

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (7)$$

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Eoption_2Eoption\ A0) \quad (8)$$

Let $ty_2Ellist_2Ellist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Ellist_2Ellist\ A0) \quad (9)$$

Let $c_2Ellist_2Ellist_rep : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Ellist_2Ellist_rep\ A_27a \in \\ (((ty_2Eoption_2Eoption\ A_27a)^{ty_2Enum_2Enum})^{(ty_2Ellist_2Ellist\ A_27a)}) \quad (10)$$

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \quad (11)$$

Definition 12 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \quad (12)$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \quad (13)$$

Definition 13 We define c_2Esum_2EINL to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27a.(ap\ (c_2Esum_2EABS$

Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Eoption_2Eoption_ABS\ A_27a \in \\ ((ty_2Eoption_2Eoption\ A_27a)^{(ty_2Esum_2Esum\ A_27a\ ty_2Eone_2Eone)}) \quad (14)$$

Definition 14 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. (ap (c_2Eoption_2Eoption_2ABS A_27a) x)$

Definition 15 We define $c_2Emin_2E_40$ to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (ap P x)) \text{ then } (the (\lambda x. x \in A) \wedge P)$ of type $\iota \Rightarrow \iota$.

Definition 16 We define c_2Ebool_2ECOND to be $\lambda A_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A_27a. (\lambda V2t2 \in A_27a. (ap (c_2Ebool_2ECOND A_27a) t1 t2))))$

Let $c_2Ellist_2Ellist_abs : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow c_2Ellist_2Ellist_abs A_27a \in ((ty_2Ellist_2Ellist A_27a)^{(ty_2Eoption_2Eoption A_27a)^{ty_2Enum_2Enum}}) \quad (15)$$

Definition 17 We define $c_2Ellist_2ELCONS$ to be $\lambda A_27a : \iota. \lambda V0h \in A_27a. \lambda V1t \in (ty_2Ellist_2Ellist A_27a) h$

Definition 18 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota. (\lambda V0P \in (2^{A_27a}). (ap V0P (ap (c_2Emin_2E_40 A_27a) P))))$

Definition 19 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone. x))$

Definition 20 We define c_2Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_2Esum_2EABS A_27a A_27b) e)$

Definition 21 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota. (ap (c_2Eoption_2Eoption_2ABS A_27a) (c_2Emin_2E_40 A_27a))$

Definition 22 We define $c_2Ellist_2ELNIL$ to be $\lambda A_27a : \iota. (ap (c_2Ellist_2Ellist_abs A_27a) (\lambda V0n \in ty_2Ellist_2Ellist A_27a. n))$

Definition 23 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (V0t1 V1t2))))$

Definition 24 We define $c_2Ellist_2ELFINITE$ to be $\lambda A_27a : \iota. (\lambda V0a0 \in (ty_2Ellist_2Ellist A_27a). (ap (c_2Emin_2E_40 A_27a) a0))$

Assume the following.

$$True \quad (16)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (17)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow \neg (p V0t)))) \quad (18)$$

Assume the following.

$$(\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in 2. (((p V0x) \Rightarrow (p V1y)) \wedge ((p V2z) \Rightarrow (p V3w))) \Rightarrow (((p V0x) \wedge (p V2z)) \Rightarrow ((p V1y) \wedge (p V3w)))))) \quad (19)$$

Assume the following.

$$\begin{aligned}
& (\forall V0x \in 2. (\forall V1y \in 2. (\forall V2z \in 2. (\forall V3w \in \\
& 2. (((p V0x) \Rightarrow (p V1y)) \wedge ((p V2z) \Rightarrow (p V3w))) \Rightarrow (((p V0x) \vee (p V2z)) \Rightarrow \\
& ((p V1y) \vee (p V3w)))))))))
\end{aligned} \tag{20}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a. \text{nonempty } A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1Q \in \\
& (2^{A_27a}). ((\forall V2x \in A_27a. ((p (ap V0P V2x)) \Rightarrow (p (ap V1Q V2x)))) \Rightarrow \\
& ((\exists V3x \in A_27a. (p (ap V0P V3x))) \Rightarrow (\exists V4x \in A_27a. (p (\\
& ap V1Q V4x)))))))))
\end{aligned} \tag{21}$$

Theorem 1

$$\begin{aligned}
& \forall A_27a. \text{nonempty } A_27a \Rightarrow ((p (ap (c_2Ellist_2ELFINITE A_27a) \\
& (c_2Ellist_2ELNIL A_27a))) \wedge (\forall V0h \in A_27a. (\forall V1t \in \\
& (ty_2Ellist_2Ellist A_27a). ((p (ap (c_2Ellist_2ELFINITE A_27a) \\
& V1t)) \Rightarrow (p (ap (c_2Ellist_2ELFINITE A_27a) (ap (ap (c_2Ellist_2ELCONS \\
& A_27a) V0h) V1t)))))))))
\end{aligned}$$