

thm\_2Ellist\_2Elrep\_\_ok\_\_cases  
(TMMBpwM4FdEqcoXt3XrDJAuiY24Q3FwdA6U)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2F$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_27E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2F$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 7** We define  $c\_2Enum\_2E0$  to be  $(ap c\_2Enum\_2EABS\_num c\_2Enum\_2EZERO\_REP)$ .

**Definition 8** We define  $c\_2Earithmetic\_2EZERO$  to be  $c\_2Enum\_2E0$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 9** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (6)$$

**Definition 10** We define  $c\_2Earithmetic\_2EBIT1$  to be  $\lambda V0n \in ty\_2Enum\_2Enum.(ap\ (ap\ c\_2Earithmetic$

**Definition 11** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Let  $c\_2Earithmetic\_2E\_2D : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2D \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \quad (7)$$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \quad (8)$$

**Definition 12** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \quad (9)$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \quad (10)$$

**Definition 13** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap\ (c\_2Esum\_2EABS$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Eoption\_2Eoption\ A0) \quad (11)$$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2Eoption\_ABS\ A\_27a \in ((ty\_2Eoption\_2Eoption\ A\_27a)^{(ty\_2Esum\_2Esum\ A\_27a\ ty\_2Eone\_2Eone)}) \quad (12)$$

**Definition 14** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.(ap\ (c\_2Eoption\_2Eoption$

**Definition 15** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if}\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge P\ x)$   
of type  $\iota \Rightarrow \iota$ .

**Definition 16** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A\_27a.(\lambda V2t2 \in A\_27a.($

**Definition 17** We define  $c\_2Eone\_2Eone$  to be  $(ap\ (c\_2Emin\_2E\_40\ ty\_2Eone\_2Eone)\ (\lambda V0x \in ty\_2Eone\_2Eone$

**Definition 18** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap (c\_2Esum\_2EABS$

**Definition 19** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A\_27a : \iota.(ap (c\_2Eoption\_2Eoption\_ABS A\_27a) (c$

**Definition 20** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

**Definition 21** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 22** We define  $c\_2Ellist\_2Elrep\_ok$  to be  $\lambda A\_27a : \iota.(\lambda V0a0 \in ((ty\_2Eoption\_2Eoption A\_27a)^{ty-}$

Assume the following.

$$True \tag{13}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{14}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \tag{15}$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1y \in 2.(\forall V2z \in 2.(\forall V3w \in 2.(((p V0x) \Rightarrow (p V1y)) \wedge ((p V2z) \Rightarrow (p V3w))) \Rightarrow (((p V0x) \wedge (p V2z)) \Rightarrow ((p V1y) \wedge (p V3w)))))) \tag{16}$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1y \in 2.(\forall V2z \in 2.(\forall V3w \in 2.(((p V0x) \Rightarrow (p V1y)) \wedge ((p V2z) \Rightarrow (p V3w))) \Rightarrow (((p V0x) \vee (p V2z)) \Rightarrow ((p V1y) \vee (p V3w)))))) \tag{17}$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0P \in (2^{A-27a}).(\forall V1Q \in (2^{A-27a}).((\forall V2x \in A\_27a.((p (ap V0P V2x)) \Rightarrow (p (ap V1Q V2x)))) \Rightarrow ((\exists V3x \in A\_27a.(p (ap V0P V3x))) \Rightarrow (\exists V4x \in A\_27a.(p (ap V1Q V4x)))))) \tag{18}$$

**Theorem 1**

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0a0 \in ((ty\_2Eoption\_2Eoption A\_27a)^{ty\_2Enum\_2Enum}).((p (ap (c\_2Ellist\_2Elrep\_ok A\_27a) V0a0)) \Leftrightarrow ((V0a0 = (\lambda V1n \in ty\_2Enum\_2Enum.(c\_2Eoption\_2ENONE A\_27a)) \vee (\exists V2h \in A\_27a.(\exists V3t \in ((ty\_2Eoption\_2Eoption A\_27a)^{ty\_2Enum\_2Enum}).((V0a0 = (\lambda V4n \in ty\_2Enum\_2Enum.(ap (ap (ap (c\_2Ebool\_2ECOND (ty\_2Eoption\_2Eoption A\_27a)) (ap (ap (c\_2Emin\_2E\_3D ty\_2Enum\_2Enum) V4n) c\_2Enum\_2E0)) (ap (c\_2Eoption\_2ESOME A\_27a) V2h)) (ap V3t (ap (ap c\_2Earithmetic\_2E\_2D V4n) (ap c\_2Earithmetic\_2ENUMERAL (ap c\_2Earithmetic\_2EBIT1 c\_2Earithmetic\_2EZERO)))))) \wedge (p (ap (c\_2Ellist\_2Elrep\_ok A\_27a) V3t)))))) \tag{19}$$