

thm\_2Emarker\_2Emove\_\_right\_\_disj  
(TMXfqfMRvMY69Bakot32cqawLaaeYY5f3Bs)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 7** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 8** We define  $c\_2Emarker\_2Estmarker$  to be  $\lambda A\_27a : \iota.\lambda V0x \in A\_27a.V0x$ .

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (2)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (3)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(\forall V2C \in 2.(((p V0A) \vee (p V1B) \vee (p V2C)) \Leftrightarrow (((p V0A) \vee (p V1B)) \vee (p V2C)))))) \quad (4)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))))) \quad (5)$$

**Theorem 1**

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2m \in 2.((((p (ap (c\_2Emarker\_2Estmarker 2) V2m)) \vee (p V0p)) \Leftrightarrow ((p V0p) \vee (p (ap (c\_2Emarker\_2Estmarker 2) V2m)))))) \wedge (((p V0p) \vee ((p V1q) \vee (p (ap (c\_2Emarker\_2Estmarker 2) V2m)))))) \Leftrightarrow (((p V0p) \vee (p V1q)) \vee (p (ap (c\_2Emarker\_2Estmarker 2) V2m)))))) \wedge (((p V0p) \vee (p (ap (c\_2Emarker\_2Estmarker 2) V2m))) \vee (p V1q)) \Leftrightarrow (((p V0p) \vee (p V1q)) \vee (p (ap (c\_2Emarker\_2Estmarker 2) V2m)))))))))$$