

thm_2Emetric_2EMETRIC__SAME
(TMUN6LxkArc6pNqxwbFGgR4b9NLMoZXq9SR)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2))) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x)$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 3 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $ty_2Erealx_2Ereal : \iota$ be given. Assume the following.

$$nonempty\ ty_2Erealx_2Ereal \tag{4}$$

Let $c_2Ereal_2Ereal_of_num : \iota$ be given. Assume the following.

$$c_2Ereal_2Ereal_of_num \in (ty_2Erealx_2Ereal^{ty_2Enum_2Enum}) \tag{5}$$

Definition 4 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})))$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2)) (\lambda V2t \in 2.$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Epair_2Eprod\ A0\ A1) \quad (6)$$

Let $c_2Epair_2EABS_prod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epair_2EABS_prod\ A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ A_27b)^{(2^{A_27b})^{A_27a}}) \quad (7)$$

Definition 7 We define $c_2Epair_2E_2C$ to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0x \in A_27a. \lambda V1y \in A_27b. (ap\ (c_2E$

Let $ty_2Emetric_2Emetric : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Emetric_2Emetric\ A0) \quad (8)$$

Let $c_2Emetric_2Edist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Emetric_2Edist\ A_27a \in ((ty_2Erealax_2Ereal^{(ty_2Epair_2Eprod\ A_27a\ A_27a)})) \quad (9)$$

Assume the following.

$$True \quad (10)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (11)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0m \in (ty_2Emetric_2Emetric\ A_27a). (\forall V1x \in A_27a. (\forall V2y \in A_27a. (((ap\ (ap\ (c_2Emetric_2Edist\ A_27a)\ V0m)\ (ap\ (ap\ (c_2Epair_2E_2C\ A_27a\ A_27a)\ V1x)\ V2y)) = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0)) \Leftrightarrow (V1x = V2y)))))) \quad (12)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0m \in (ty_2Emetric_2Emetric\ A_27a). (\forall V1x \in A_27a. ((ap\ (ap\ (c_2Emetric_2Edist\ A_27a)\ V0m)\ (ap\ (ap\ (c_2Epair_2E_2C\ A_27a\ A_27a)\ V1x)\ V1x)) = (ap\ c_2Ereal_2Ereal_of_num\ c_2Enum_2E0))))))$$