

thm_2Enum_2EINV__SUC
(TMQZtm1mGc76v9Xv4me4ojGMMcQVWZ1Sena)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ **then** $(the (\lambda x.x \in A \wedge p (ap P x)))$ of type $\iota \Rightarrow \iota$.

Definition 6 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a P))))$

Definition 7 We define $c_2Ebool_2E_ONTO$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0f \in (A_27b^{A_27a}).(ap (c_2Ebool_2E_21 2) (ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap (c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F 2) (ap (ap (c_2Emin_2E_3D (2^{2^2}))$

Definition 10 We define $c_2Ebool_2E_ONE_ONE$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.(\lambda V0f \in (A_27b^{A_27a}).(ap (c_2Ebool_2E_21 2) (ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{2}$$

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t) (ap (ap (c_2Emin_2E_3D (2^{2^2}))$

Definition 12 We define $c_2Enum_2EIS_NUM_REP$ to be $\lambda V0m \in \omega.(ap (c_2Ebool_2E_21 (2^{\omega})) (ap (ap (c_2Emin_2E_3D (2^{2^2}))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \quad (3)$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (5)$$

Definition 13 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (9)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow (\forall V0f \in (A.27b^{A.27a}).((p\ (ap\ (c_2Ebool_2EONE_ONE\ A.27a\ A.27b)\ V0f)) \Leftrightarrow (\forall V1x1 \in A.27a.(\forall V2x2 \in A.27a.(((ap\ V0f\ V1x1) = (ap\ V0f\ V2x2)) \Rightarrow (V1x1 = V2x2)))))) \quad (10)$$

Assume the following.

$$((p\ (ap\ (c_2Ebool_2EONE_ONE\ \omega\ \omega)\ c_2Enum_2ESUC_REP)) \wedge (\neg(p\ (ap\ (c_2Ebool_2EONTO\ \omega\ \omega)\ c_2Enum_2ESUC_REP)))) \quad (11)$$

Assume the following.

$$((\forall V0a \in ty_2Enum_2Enum.((ap\ c_2Enum_2EABS_num\ (ap\ c_2Enum_2EREP_num\ V0a)) = V0a)) \wedge (\forall V1r \in \omega.((p\ (ap\ c_2Enum_2EIS_NUM_REP\ V1r)) \Leftrightarrow ((ap\ c_2Enum_2EREP_num\ (ap\ c_2Enum_2EABS_num\ V1r)) = V1r)))) \quad (12)$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.((ap\ c_2Enum_2ESUC\ V0m) = (ap\ c_2Enum_2ESUC\ V1n)) \Rightarrow (V0m = V1n))))$$