

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (5)$$

Definition 10 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (9)$$

Assume the following.

$$(\forall V0y \in \omega.(\neg(c_2Enum_2EZERO_REP = (ap\ c_2Enum_2ESUC_REP\ V0y)))) \quad (10)$$

Assume the following.

$$((\forall V0a \in ty_2Enum_2Enum.((ap\ c_2Enum_2EABS_num\ (ap\ c_2Enum_2EREP_num\ V0a)) = V0a)) \wedge (\forall V1r \in \omega.((p\ (ap\ c_2Enum_2EIS_NUM_REP\ V1r)) \Leftrightarrow ((ap\ c_2Enum_2EREP_num\ (ap\ c_2Enum_2EABS_num\ V1r)) = V1r)))) \quad (11)$$

Theorem 1

$$(\forall V0n \in ty_2Enum_2Enum.(\neg((ap\ c_2Enum_2ESUC\ V0n) = c_2Enum_2E0)))$$