

thm_2Enumeral_2Enumeral__pre (TMMb- BjY4Ja63KP9H8ZV9fm6KvnbVPQtuxXh)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \tag{1}$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{2}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{3}$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Definition 3 We define $c_2Earithmetic_2EZERO$ to be c_2Enum_2E0 .

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{4}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{5}$$

Definition 4 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap\ (ap\ (c_2Emin_2E_3D\ (2^{A-27a}))\ (\lambda V1x \in 2.V1x))\ (\lambda V2x \in 2.V2x)))$

Definition 6 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap\ c_2Enum_2EABS_num\ (ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))))$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \tag{6}$$

Definition 7 We define `c_2Earithmic_2EBIT1` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT1 V0n))$.

Definition 8 We define `c_2Earithmic_2EBIT2` to be $\lambda V0n \in ty_2Enum_2Enum.(ap (ap c_2Earithmic_2EBIT2 V0n))$.

Definition 9 We define `c_2Ebool_2EF` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 11 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 12 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$.

Definition 13 We define `c_2Ebool_2ECOND` to be $\lambda A.\lambda a : \iota.(\lambda V0t \in 2.(\lambda V1t1 \in A.\lambda V2t2 \in A.\lambda V3t3 \in A.(ap (c_2Ebool_2E_21 2) (ap (ap (ap (c_2Ebool_2E_21 2) V0t) V1t1) V2t2) V3t3))))$.

Definition 14 We define `c_2Eprim_rec_2EPRE` to be $\lambda V0m \in ty_2Enum_2Enum.(ap (ap (ap (c_2Ebool_2E_21 2) V0m) V1n) V2p))$.

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & ((ap (ap c_2Earithmic_2E_2B c_2Enum_2E0) V0m) = V0m) \wedge (((ap (\\ & ap c_2Earithmic_2E_2B V0m) c_2Enum_2E0) = V0m) \wedge (((ap (ap c_2Earithmic_2E_2B \\ & (ap c_2Enum_2ESUC V0m)) V1n) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmic_2E_2B \\ & V0m) V1n)))) \wedge ((ap (ap c_2Earithmic_2E_2B V0m) (ap c_2Enum_2ESUC \\ & V1n)) = (ap c_2Enum_2ESUC (ap (ap c_2Earithmic_2E_2B V0m) V1n)))))))) \\ & \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0m \in ty_2Enum_2Enum.(\forall V1n \in ty_2Enum_2Enum.(\\ & \forall V2p \in ty_2Enum_2Enum.((ap (ap c_2Earithmic_2E_2B V0m) \\ & (ap (ap c_2Earithmic_2E_2B V1n) V2p)) = (ap (ap c_2Earithmic_2E_2B \\ & (ap (ap c_2Earithmic_2E_2B V0m) V1n)) V2p)))) \\ & \end{aligned} \tag{8}$$

Assume the following.

$$True \tag{9}$$

Assume the following.

$$\forall A.\lambda a.nonempty A.\lambda a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A.(ap (ap c_2Earithmic_2E_2B V0t) V1x) = V1x) \Leftrightarrow (ap (ap c_2Earithmic_2E_2B V0t) V1x) = V1x))) \tag{10}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \\ & \end{aligned} \tag{11}$$

Assume the following.

$$\forall A.\lambda a.nonempty A.\lambda a \Rightarrow (\forall V0x \in A.(V0x = V0x) \Leftrightarrow True) \tag{12}$$

Assume the following.

$$\begin{aligned}
&(((ap\ c_2Eprim_rec_2EPRE\ c_2Enum_2E0) = c_2Enum_2E0) \wedge (\forall V0m \in \\
&ty_2Enum_2Enum.((ap\ c_2Eprim_rec_2EPRE\ (ap\ c_2Enum_2ESUC\ V0m)) = \\
&V0m))) \\
& \hspace{15em} (13)
\end{aligned}$$

Theorem 1

$$\begin{aligned}
&(((ap\ c_2Eprim_rec_2EPRE\ c_2Earithmetic_2EZERO) = c_2Earithmetic_2EZERO) \wedge \\
&(((ap\ c_2Eprim_rec_2EPRE\ (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO)) = \\
& \quad c_2Earithmetic_2EZERO) \wedge ((\forall V0n \in ty_2Enum_2Enum.((ap \\
& \quad c_2Eprim_rec_2EPRE\ (ap\ c_2Earithmetic_2EBIT1\ (ap\ c_2Earithmetic_2EBIT1 \\
& \quad V0n))) = (ap\ c_2Earithmetic_2EBIT2\ (ap\ c_2Eprim_rec_2EPRE\ (ap \\
& \quad c_2Earithmetic_2EBIT1\ V0n)))))) \wedge ((\forall V1n \in ty_2Enum_2Enum. \\
& ((ap\ c_2Eprim_rec_2EPRE\ (ap\ c_2Earithmetic_2EBIT1\ (ap\ c_2Earithmetic_2EBIT2 \\
& \quad V1n))) = (ap\ c_2Earithmetic_2EBIT2\ (ap\ c_2Earithmetic_2EBIT1 \\
& \quad V1n)))))) \wedge (\forall V2n \in ty_2Enum_2Enum.((ap\ c_2Eprim_rec_2EPRE \\
& (ap\ c_2Earithmetic_2EBIT2\ V2n)) = (ap\ c_2Earithmetic_2EBIT1\ V2n))))))
\end{aligned}$$