

thm\_2Enumeral\_bit\_2EiDUB\_\_NUMERAL  
(TMFroTjywbdCyHv4qWj2hzigJuSqQLGZC9xo)

October 26, 2020

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Earithmetic\_2E\_2B : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2B \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{2}$$

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Enumeral\_2EiDUB$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.(ap (ap c\_2Earithmetic\_2E\_2B$

**Definition 5** We define  $c\_2Earithmetic\_2ENUMERAL$  to be  $\lambda V0x \in ty\_2Enum\_2Enum.V0x$ .

Assume the following.

$$True \tag{3}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \tag{4}$$

**Theorem 1**

$$(\forall V0i \in ty\_2Enum\_2Enum.((ap\ c\_2Enumeral\_2EiDUB\ (ap\ c\_2Earithmetic\_2ENUMERAL\ V0i)) = (ap\ c\_2Earithmetic\_2ENUMERAL\ (ap\ c\_2Enumeral\_2EiDUB\ V0i))))$$