

thm_2Eoption_2ENOT__SOME__NONE (TMU- jXwDbhVFWTCFw6jYhr4Z25pZvdSkWj6q)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow P \Rightarrow Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a})))$

Definition 5 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Eoption_2Eoption\ A0) \tag{3}$$

Let $c_2Eoption_2Eoption_REP : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Eoption_2Eoption_REP\ A_27a \in ((ty_2Esum_2Esum\ A_27a\ ty_2Eone_2Eone)^{(ty_2Eoption_2Eoption\ A_27a)}) \tag{4}$$

Definition 6 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \tag{5}$$

Definition 7 We define c_2Esum_2EINL to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27a.(ap (c_2Esum_2EABS_27a) (c_2Eoption_2Eoption_ABS A_27a))$.
Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Eoption_2Eoption_ABS A_27a \in ((ty_2Eoption_2Eoption A_27a)^{(ty_2Esum_2Esum A_27a ty_2Eone_2Eone)}) \quad (6)$$

Definition 8 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota.\lambda V0x \in A_27a.(ap (c_2Eoption_2Eoption_ABS A_27a) (V0x))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 10 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone.V0x))$

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_7E))$

Definition 12 We define c_2Esum_2EINR to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27b.(ap (c_2Esum_2EABS_27a) (V0e))$

Definition 13 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota.(ap (c_2Eoption_2Eoption_ABS A_27a) (V0))$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (8)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \quad (9)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.(((\forall V1x \in A_27a.(p V0t) \Leftrightarrow (p V0t)))))) \quad (10)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \quad (11)$$

Assume the following.

$$((\forall V0t \in 2.(((\neg (\neg (p V0t))) \Leftrightarrow (p V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)))))) \quad (12)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow ((\forall V0a \in (ty_2Eoption_2Eoption \\
& A.27a).(ap\ (c_2Eoption_2Eoption_ABS\ A.27a)\ (ap\ (c_2Eoption_2Eoption_REP \\
& A.27a)\ V0a)) = V0a)) \wedge (\forall V1r \in (ty_2Esum_2Esum\ A.27a\ ty_2Eone_2Eone). \\
& ((p\ (ap\ (\lambda V2x \in (ty_2Esum_2Esum\ A.27a\ ty_2Eone_2Eone).c_2Ebool_2ET) \\
& V1r)) \Leftrightarrow ((ap\ (c_2Eoption_2Eoption_REP\ A.27a)\ (ap\ (c_2Eoption_2Eoption_ABS \\
& A.27a)\ V1r)) = V1r))))
\end{aligned} \tag{14}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow \forall A.27b.nonempty\ A.27b \Rightarrow (\\
& \forall V0v1 \in A.27a.(\forall V1v2 \in A.27b.(\neg((ap\ (c_2Esum_2EINR \\
& A.27a\ A.27b)\ V1v2) = (ap\ (c_2Esum_2EINL\ A.27a\ A.27b)\ V0v1))))))
\end{aligned} \tag{15}$$

Theorem 1

$$\begin{aligned}
& \forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a.(\neg((ap\ (c_2Eoption_2ESOME \\
& A.27a)\ V0x) = (c_2Eoption_2ENONE\ A.27a))))
\end{aligned}$$