

# thm\_2Eoption\_2EOPTION\_\_CHOICE\_\_NONE (TMacJeQC7wKmvzww1n9ybKgjjKvgsBsb5cL)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p x)$ ) of type  $\iota \Rightarrow \iota$ .

**Definition 4** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40 A\_27a P))))$

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}) P) P)))$

**Definition 7** We define  $c\_2Ebool\_2E\_5C\_2E\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \tag{1}$$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \tag{2}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{3}$$

**Definition 9** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b) V0e)$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Eoption\_2Eoption\ A0) \quad (4)$$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow c\_2Eoption\_2Eoption\_ABS\ A.27a \in ((ty\_2Eoption\_2Eoption\ A.27a)^{(ty\_2Esum\_2Esum\ A.27a\ ty\_2Eone\_2Eone)}) \quad (5)$$

**Definition 10** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A.27a : \iota. \lambda V0x \in A.27a. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A.27a)\ V0x)$

**Definition 11** We define  $c\_2Eone\_2Eone$  to be  $(ap\ (c\_2Emin\_2E\_40\ ty\_2Eone\_2Eone)\ (\lambda V0x \in ty\_2Eone\_2Eone.\ V0x))$

**Definition 12** We define  $c\_2Ebool\_2EF$  to be  $(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V0t \in 2.V0t))$ .

**Definition 13** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2. (ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_7E))$

**Definition 14** We define  $c\_2Esum\_2EINR$  to be  $\lambda A.27a : \iota. \lambda A.27b : \iota. \lambda V0e \in A.27b. (ap\ (c\_2Esum\_2EABS\ A.27a)\ V0e)$

**Definition 15** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A.27a : \iota. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A.27a)\ (c\_2Eoption\_2ENONE\ A.27a))$

Let  $c\_2Eoption\_2EOPTION\_CHOICE : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow c\_2Eoption\_2EOPTION\_CHOICE\ A.27a \in (((ty\_2Eoption\_2Eoption\ A.27a)^{(ty\_2Eoption\_2Eoption\ A.27a)})^{(ty\_2Eoption\_2Eoption\ A.27a)}) \quad (6)$$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a. ((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0opt \in (ty\_2Eoption\_2Eoption\ A.27a). ((V0opt = (c\_2Eoption\_2ENONE\ A.27a)) \vee (\exists V1x \in A.27a. (V0opt = (ap\ (c\_2Eoption\_2ESOME\ A.27a)\ V1x)))))) \quad (9)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow (\forall V0x \in A.27a. (\forall V1y \in A.27a. (((ap\ (c\_2Eoption\_2ESOME\ A.27a)\ V0x) = (ap\ (c\_2Eoption\_2ESOME\ A.27a)\ V1y)) \Leftrightarrow (V0x = V1y)))) \quad (10)$$

Assume the following.

$$\forall A.27a.nonempty\ A.27a \Rightarrow ((\forall V0m2 \in (ty\_2Eoption\_2Eoption\ A.27a). ((ap\ (ap\ (c\_2Eoption\_2EOPTION\_CHOICE\ A.27a)\ (c\_2Eoption\_2ENONE\ A.27a))\ V0m2) = V0m2)) \wedge (\forall V1x \in A.27a. (\forall V2m2 \in (ty\_2Eoption\_2Eoption\ A.27a). ((ap\ (ap\ (c\_2Eoption\_2EOPTION\_CHOICE\ A.27a)\ (ap\ (c\_2Eoption\_2ESOME\ A.27a)\ V1x))\ V2m2) = (ap\ (c\_2Eoption\_2ESOME\ A.27a)\ V1x)))))) \quad (11)$$

**Theorem 1**

$\forall A_{27a}.nonempty\ A_{27a} \Rightarrow (\forall V0m1 \in (ty\_2Eoption\_2Eoption$   
 $A_{27a}).((ap\ (ap\ (c\_2Eoption\_2EOPTION\_CHOICE\ A_{27a})\ V0m1)\ (c\_2Eoption\_2ENONE$   
 $A_{27a})) = V0m1))$