

thm_2Eoption_2Eoption_CASES (TMPt- DVG5UPwFhNzeAHJ8E1dpQWpVTA1HNYK)

October 26, 2020

Let $ty_2Eone_2Eone : \iota$ be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Definition 1 We define c_2Emin_2E3D to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Emin_2E3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 3 We define c_2Ebool_2E2T to be $(ap (ap (c_2Emin_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define c_2Ebool_2E21 to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E3D (2^{A_27a})))$

Definition 5 We define $c_2Ebool_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E21 2) (\lambda V2t \in 2.V2t)))$

Let $ty_2Esum_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let $c_2Esum_2EABS_sum : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Esum_2EABS_sum\ A_27a\ A_27b \in ((ty_2Esum_2Esum\ A_27a\ A_27b)^{((2^{A_27b})^{A_27a})^2}) \tag{3}$$

Definition 6 We define c_2Esum_2EINL to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27a.(ap (c_2Esum_2EABS_sum$

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty_2Eoption_2Eoption\ A0) \tag{4}$$

Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow c_2Eoption_2Eoption_ABS\ A_27a \in ((ty_2Eoption_2Eoption\ A_27a)^{(ty_2Esum_2Esum\ A_27a\ ty_2Eone_2Eone)}) \tag{5}$$

Definition 7 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota.\lambda V0x \in A_27a.(ap (c_2Eoption_2Eoption_2ABS A_27a) V0x)$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.\text{if } (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a) V0P)))$

Definition 10 We define c_2Eone_2Eone to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone V0x))$

Definition 11 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 12 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 13 We define c_2Esum_2EINR to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0e \in A_27b.(ap (c_2Esum_2EABS A_27a A_27b) V0e)$

Definition 14 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota.(ap (c_2Eoption_2Eoption_2ABS A_27a) (the (empty)))$

Definition 15 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t) V1t2) V0t1))$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \vee (p V1B)) \Leftrightarrow ((p V1B) \vee (p V0A)))))) \quad (6)$$

Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0opt \in (ty_2Eoption_2Eoption A_27a).((V0opt = (c_2Eoption_2ENONE A_27a)) \vee (\exists V1x \in A_27a.(V0opt = (ap (c_2Eoption_2ESOME A_27a) V1x)))))) \quad (7)$$

Theorem 1

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0opt \in (ty_2Eoption_2Eoption A_27a).((\exists V1x \in A_27a.(V0opt = (ap (c_2Eoption_2ESOME A_27a) V1x))) \vee (V0opt = (c_2Eoption_2ENONE A_27a))))))$$