

thm_2Eoption_2Eoption__case__SOME_ID
(TMcirUeLs2AhZUvRrXGRqnv2Nn8nHwR78qb)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Emin_2E_40` to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p x))$ of type $\iota \Rightarrow \iota$.

Definition 4 We define `c_2Ebool_2E_3F` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap V0P (ap (c_2Emin_2E_40 A_{27a} P))))$

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap (ap (c_2Emin_2E_3D (2^{A_{27a}}) P)) V0P)))$

Definition 7 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Let `ty_2Eone_2Eone` : ι be given. Assume the following.

$$nonempty\ ty_2Eone_2Eone \tag{1}$$

Definition 8 We define `c_2Eone_2Eone` to be $(ap (c_2Emin_2E_40 ty_2Eone_2Eone) (\lambda V0x \in ty_2Eone_2Eone.V0x))$

Definition 9 We define `c_2Ebool_2E_2F` to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2F))$

Definition 11 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t))))$

Let `ty_2Esum_2Esum` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty_2Esum_2Esum\ A0\ A1) \tag{2}$$

Let `c_2Esum_2EABS__sum` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_{27a}.nonempty\ A_{27a} \Rightarrow \forall A_{27b}.nonempty\ A_{27b} \Rightarrow c_2Esum_2EABS_sum\ A_{27a}\ A_{27b} \in ((ty_2Esum_2Esum\ A_{27a}\ A_{27b})^{((2^{A_{27b}})^{A_{27a}})^2}) \tag{3}$$

Definition 12 We define c_2Esum_2EINR to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27b. (ap (c_2Esum_2EABS$
Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. nonempty A0 \Rightarrow nonempty (ty_2Eoption_2Eoption A0) \quad (4)$$

Let $c_2Eoption_2Eoption_ABS : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow c_2Eoption_2Eoption_ABS A_27a \in ((ty_2Eoption_2Eoption A_27a)^{(ty_2Esum_2Esum A_27a ty_2Eone_2Eone)}) \quad (5)$$

Definition 13 We define $c_2Eoption_2ENONE$ to be $\lambda A_27a : \iota. (ap (c_2Eoption_2Eoption_ABS A_27a) (c_2Eoption_2Eoption_ABS A_27a))$

Let $c_2Eoption_2Eoption_CASE : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow c_2Eoption_2Eoption_CASE A_27a A_27b \in (((A_27b)^{(A_27b^{A_27a}})^{A_27b})^{(ty_2Eoption_2Eoption A_27a)}) \quad (6)$$

Definition 14 We define c_2Esum_2EINL to be $\lambda A_27a : \iota. \lambda A_27b : \iota. \lambda V0e \in A_27a. (ap (c_2Esum_2EABS$

Definition 15 We define $c_2Eoption_2ESOME$ to be $\lambda A_27a : \iota. \lambda V0x \in A_27a. (ap (c_2Eoption_2Eoption_ABS$

Assume the following.

$$True \quad (7)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0opt \in (ty_2Eoption_2Eoption A_27a). ((V0opt = (c_2Eoption_2ENONE A_27a) \vee (\exists V1x \in A_27a. (V0opt = (ap (c_2Eoption_2ESOME A_27a) V1x)))))) \quad (9)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow \forall A_27b. nonempty A_27b \Rightarrow ((\forall V0v \in A_27b. (\forall V1f \in (A_27b^{A_27a}). ((ap (ap (ap (c_2Eoption_2Eoption_CASE A_27a A_27b) (c_2Eoption_2ENONE A_27a)) V0v) V1f) = V0v))) \wedge (\forall V2x \in A_27a. (\forall V3v \in A_27b. (\forall V4f \in (A_27b^{A_27a}). ((ap (ap (ap (c_2Eoption_2Eoption_CASE A_27a A_27b) (ap (c_2Eoption_2ESOME A_27a) V2x)) V3v) V4f) = (ap V4f V2x))))))) \quad (10)$$

Assume the following.

$$\forall A_27a. nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in A_27a. (((ap (c_2Eoption_2ESOME A_27a) V0x) = (ap (c_2Eoption_2ESOME A_27a) V1y)) \Leftrightarrow (V0x = V1y)))) \quad (11)$$

Theorem 1

$$\forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0x \in (\text{ty_2Eoption_2Eoption } A_{27a}). ((\text{ap } (\text{ap } (\text{ap } (\text{c_2Eoption_2Eoption_CASE } A_{27a} (\text{ty_2Eoption_2Eoption } A_{27a})) V0x) V0x) (\text{c_2Eoption_2ESOME } A_{27a})) = V0x))$$