

thm_2Epath_2Edrop_def_compute
 (TMTBqWZnRrfafnKD9s8pvT8jqzeAuS1Anbq)

October 26, 2020

Let $c_2Enum_2EZERO_REP : \iota$ be given. Assume the following.

$$c_2Enum_2EZERO_REP \in \omega \quad (1)$$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$\text{nonempty } ty_2Enum_2Enum \quad (2)$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \quad (3)$$

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A. \lambda x \in A. \lambda y \in A. inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Enum_2E0 to be $(ap\ c_2Enum_2EABS_num\ c_2Enum_2EZERO_REP)$.

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \quad (4)$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \quad (5)$$

Definition 3 We define c_2Ebool_2ET to be $(ap\ (ap\ (c_2Emin_2E_3D\ (2^2))\ (\lambda V0x \in 2.V0x))\ (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A. \lambda P \in (2^{A \rightarrow 2}). (ap\ (ap\ (c_2Emin_2E_3D\ (2^{A \rightarrow 2}))\ (\lambda V0P \in 2^{A \rightarrow 2})))$

Definition 5 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum. (ap\ c_2Enum_2EABS_num\ m)$

Let $c_2Earithmetic_2E_2B : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2B \in ((ty_2Enum_2Enum^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}) \quad (6)$$

Definition 6 We define $c_2Earithmetic_2EBIT2$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earithmetic_2EBIT2\ n)\ V)$

Definition 7 We define $\text{c_2Earithmetic_2EZERO}$ to be c_2Enum_2E0 .

Let $c_2Earithmetic_2E_2D : \iota$ be given. Assume the following.

$$c_2Earithmetic_2E_2D \in ((ty_2Enum_2Enum^{ty_2Enum^{ty_2Enum_2Enum}})^{ty_2Enum_2Enum}) \quad (7)$$

Definition 8 We define $c_2Earthmetic_2EBIT1$ to be $\lambda V0n \in ty_2Enum_2Enum.(ap\ (ap\ c_2Earthmetic_2EBIT1\ n)\ V)$

Definition 9 We define `c_2Earthmetic_2ENUMERAL` to be $\lambda V0x \in ty_2Enum_2Enum. V0x$.

Definition 10 We define $c_2 \in \text{min_3D_3D_3E}$ to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o} (p \rightarrow p \cdot Q)$ of type ι .

Let $ty_2Epath_2Epath : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.\text{nonempty } A0 \Rightarrow \forall A1.\text{nonempty } A1 \Rightarrow \text{nonempty}(\text{ty_2Epath_2Epath } A0 \ A1)$$

Let $c_2Epath_2Etail : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow c_2\text{Epath}_2\text{Etail}_A_27a \ A_27b \in ((\text{ty}_2\text{Epath}_2\text{Epath } A_27a \ A_27b)^{(\text{ty}_2\text{Epath}_2\text{Epath } A_27a \ A_27b)}) \quad (9)$$

Let $c_2Epath_2Edrop : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A_{27a}.nonempty\ A_{27a} \Rightarrow \forall A_{27b}.nonempty\ A_{27b} \Rightarrow c_2Epath_2Edrop \\ & A_{27a}\ A_{27b} \in (((ty_2Epath_2Epath\ A_{27a}\ A_{27b})^{(ty_2Epath_2Epath\ A_{27a}\ A_{27b})})^{ty_2Enum_2Enum}) \end{aligned} \quad (10)$$

Definition 11 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Assume the following.

$\forall A_27a.\text{nonempty } A_27a \Rightarrow (\forall V0f \in ((A_27a^{ty_2Enum_2Enum})^{ty_2Enum_2Enum}).$
 $(\forall V1g \in (A_27a^{ty_2Enum_2Enum})).(\forall V2n \in ty_2Enum_2Enum.$
 $((ap\ V1g\ (ap\ c_2Enum_2ESUC\ V2n)) = (ap\ (ap\ V0f\ V2n)\ (ap\ c_2Enum_2ESUC\ V2n))) \Leftrightarrow (\forall V3n \in ty_2Enum_2Enum.((ap\ V1g\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ V3n))) = (ap\ (ap\ V0f\ (ap\ (ap\ c_2Earithmetic_2E_2D\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ V3n))))$
 $(ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ c_2Earithmetic_2EZERO)))) \wedge$
 $(\forall V4n \in ty_2Enum_2Enum.((ap\ V1g\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT2\ V4n))) = (ap\ (ap\ V0f\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT1\ V4n)))\ (ap\ c_2Earithmetic_2ENUMERAL\ (ap\ c_2Earithmetic_2EBIT2\ V4n)))))))$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow (\\ & (\forall V0p \in (\text{ty_2Epath_2Epath } A_27a _A_27b).((\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) \text{ c_2Enum_2E0}) V0p) = V0p)) \wedge (\forall V1n \in \text{ty_2Enum_2Enum}. \\ & (\forall V2p \in (\text{ty_2Epath_2Epath } A_27a _A_27b).((\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) (\text{ap } \text{c_2Enum_2ESUC } V1n)) V2p) = (\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) V1n) (\text{ap } (\text{c_2Epath_2Etal } A_27a _A_27b) V2p))))))) \end{aligned} \quad (13)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.\text{nonempty } A_27a \Rightarrow \forall A_27b.\text{nonempty } A_27b \Rightarrow (\\ & (\forall V0p \in (\text{ty_2Epath_2Epath } A_27a _A_27b).((\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) \text{ c_2Enum_2E0}) V0p) = V0p)) \wedge ((\forall V1n \in \text{ty_2Enum_2Enum}. \\ & (\forall V2p \in (\text{ty_2Epath_2Epath } A_27a _A_27b).((\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) (\text{ap } \text{c_2Earithmetic_2ENUMERAL } (\text{ap } \text{c_2Earithmetic_2EBIT1 } \\ & V1n))) V2p) = (\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } A_27a _A_27b) (\text{ap } (\text{ap } \text{c_2Earithmetic_2E_2D } \\ & (\text{ap } \text{c_2Earithmetic_2ENUMERAL } (\text{ap } \text{c_2Earithmetic_2EBIT1 } V1n)))) \\ & (\text{ap } \text{c_2Earithmetic_2ENUMERAL } (\text{ap } \text{c_2Earithmetic_2EBIT1 } \text{c_2Earithmetic_2EZERO}))) \\ & (\text{ap } (\text{c_2Epath_2Etal } A_27a _A_27b) V2p)))))) \wedge (\forall V3n \in \text{ty_2Enum_2Enum}. \\ & (\forall V4p \in (\text{ty_2Epath_2Epath } A_27a _A_27b).((\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } \\ & A_27a _A_27b) (\text{ap } \text{c_2Earithmetic_2ENUMERAL } (\text{ap } \text{c_2Earithmetic_2EBIT2 } \\ & V3n))) V4p) = (\text{ap } (\text{ap } (\text{c_2Epath_2Edrop } A_27a _A_27b) (\text{ap } \text{c_2Earithmetic_2ENUMERAL } \\ & (\text{ap } \text{c_2Earithmetic_2EBIT1 } V3n))) (\text{ap } (\text{c_2Epath_2Etal } A_27a _A_27b) \\ & V4p))))))) \end{aligned}$$