

thm_2Epath_2EfromPath_onto (TMUK6arh6gEBPwEsmw5ij3qjTVh7UngK7Wu)

October 26, 2020

Definition 1 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \ x)) \text{ then (the } (\lambda x. x \in A \wedge p \ x)) \text{ of type } \iota \Rightarrow \iota$.

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P \text{ (ap (c_2Emin_2E_40 } A \ P))$

Definition 4 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 5 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap (c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 6 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap (c_2Emin_2E_3D } (2^{A-27a}) \ P))$

Definition 7 We define `c_2Ebool_2E_2F` to be $(\text{ap (c_2Ebool_2E_21 } 2) (\lambda V0t \in 2. V0t))$.

Definition 8 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap c_2Emin_2E_3D_3D_3E } V0t) \text{ c_2Ebool_2E_2F } V0t))$

Let $ty_2Ellist_2Ellist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \text{nonempty } (ty_2Ellist_2Ellist \ A0) \quad (1)$$

Let $ty_2Epair_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (ty_2Epair_2Eprod \ A0 \ A1) \quad (2)$$

Let $ty_2Epath_2Epath : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (ty_2Epath_2Epath \ A0 \ A1) \quad (3)$$

Let $c_2Epath_2EfromPath : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epath_2EfromPath \\ & A_27a\ A_27b \in ((ty_2Epair_2Eprod\ A_27a\ (ty_2Ellist_2Ellist\ (ty_2Epair_2Eprod \\ & \quad A_27b\ A_27a)))^{(ty_2Epath_2Epath\ A_27a\ A_27b)}) \end{aligned} \quad (4)$$

Let $c_2Epath_2EtoPath : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2Epath_2EtoPath \\ & A_27a\ A_27b \in ((ty_2Epath_2Epath\ A_27a\ A_27b)^{(ty_2Epair_2Eprod\ A_27a\ (ty_2Ellist_2Ellist\ (ty_2Epair_2Eprod \end{aligned} \quad (5)$$

Definition 9 We define $c_2Ebool_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E21\ 2)\ (\lambda V2t \in 2$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p \\ & \quad V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(\quad (7) \\ & \quad p\ V0t)))))) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & (\forall V0a \in (ty_2Epath_2Epath\ A_27a\ A_27b).((ap\ (c_2Epath_2EtoPath \\ & \quad A_27a\ A_27b)\ (ap\ (c_2Epath_2EfromPath\ A_27a\ A_27b)\ V0a)) = V0a)) \wedge \\ & (\forall V1r \in (ty_2Epair_2Eprod\ A_27a\ (ty_2Ellist_2Ellist\ (ty_2Epair_2Eprod \\ & \quad A_27b\ A_27a))).((p\ (ap\ (\lambda V2x \in (ty_2Epair_2Eprod\ A_27a\ (ty_2Ellist_2Ellist \\ & \quad (ty_2Epair_2Eprod\ A_27b\ A_27a))).c_2Ebool_2ET)\ V1r)) \Leftrightarrow ((ap\ (\\ & \quad c_2Epath_2EfromPath\ A_27a\ A_27b)\ (ap\ (c_2Epath_2EtoPath\ A_27a \\ & \quad \quad A_27b)\ V1r)) = V1r)))) \end{aligned} \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & \forall V0r \in (ty_2Epair_2Eprod\ A_27a\ (ty_2Ellist_2Ellist\ (ty_2Epair_2Eprod \\ & \quad A_27b\ A_27a))).(\exists V1a \in (ty_2Epath_2Epath\ A_27a\ A_27b). \\ & \quad (V0r = (ap\ (c_2Epath_2EfromPath\ A_27a\ A_27b)\ V1a)))) \end{aligned}$$