

thm_2EpatternMatches_2EPMATCH__EQUIV__ROWS__MATCH (TMQVTqA2BQ3GFvfXqMoCtC3tBzgjnH2SZxG)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p \Rightarrow q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2EF$

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Eoption_2Eoption A0) \quad (1)$$

Let $c_2Eoption_2EIS_SOME : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Eoption_2EIS_SOME A_27a \in (2^{(ty_2Eoption_2Eoption A_27a)}) \quad (2)$$

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (3)$$

Let $c_2Elist_2ELIST_TO_SET : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2ELIST_TO_SET A_27a \in ((2^{A_27a})^{(ty_2Elist_2Elist A_27a)}) \quad (4)$$

Definition 7 We define c_2Ebool_2EIN to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 8 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 9 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A$

Let $c_2EpatternMatches_2EPMATCH : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow c_2EpatternMatches_2EPMATCH A_27a A_27b \in ((A_27a^{(ty_2Elist_2Elist ((ty_2Eoption_2Eoption A_27a)^{A_27b})})^{A_27b}) \quad (5)$$

Definition 10 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 11 We define $c_2EpatternMatches_2EPMATCH_EQUIV_ROWS$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0v \in A_27a.\lambda V1rows1 \in (ty_2Elist_2Elist ((ty_2Eoption_2Eoption A_27b)^{A_27a})$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2.(\forall V1x_27 \in 2.(\forall V2y \in 2.(\forall V3y_27 \in \\ & 2.(((p V0x) \Leftrightarrow (p V1x_27)) \wedge ((p V1x_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_27) \Rightarrow (p V3y_27)))))) \end{aligned} \quad (11)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow \forall A_27b.nonempty A_27b \Rightarrow (\\ & \forall V0v \in A_27a.(\forall V1rows1 \in (ty_2Elist_2Elist ((ty_2Eoption_2Eoption \\ & A_27b)^{A_27a})).(\forall V2rows2 \in (ty_2Elist_2Elist ((ty_2Eoption_2Eoption \\ & A_27b)^{A_27a})).((p (ap (ap (ap (c_2EpatternMatches_2EPMATCH_EQUIV_ROWS \\ & A_27a A_27b) V0v) V1rows1) V2rows2)) \Rightarrow ((ap (ap (c_2EpatternMatches_2EPMATCH \\ & A_27b A_27a) V0v) V1rows1) = (ap (ap (c_2EpatternMatches_2EPMATCH \\ & A_27b A_27a) V0v) V2rows2)))))) \end{aligned}$$