

thm_2EpatternMatches_2EPMATCH__EQUIV__ROWS__is__equiv__
(TMT7JBhfJTDcZ1eCKM55UHyiT6WUBVX6CR8)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E21 2) (\lambda V0t \in 2.V0t))$.

Let $ty_2Eoption_2Eoption : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Eoption_2Eoption A0) \quad (1)$$

Let $c_2Eoption_2Eeis_some : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Eoption_2Eeis_some A_27a \in (2^{(ty_2Eoption_2Eoption A_27a)}) \quad (2)$$

Let $ty_2Elist_2Elist : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty_2Elist_2Elist A0) \quad (3)$$

Let $c_2Elist_2Elist_to_set : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow c_2Elist_2Elist_to_set A_27a \in ((2^{A_27a})^{(ty_2Elist_2Elist A_27a)}) \quad (4)$$

Definition 5 We define $c_2Ebool_2E_2EIN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 6 We define $c_2Emin_2E_2E40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge p$ of type $\iota \Rightarrow \iota$.

Definition 7 We define $c_2Ebool_2E_2E3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_2E40 A$

Let $c_2EpatternMatches_2EPMATCH : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow c_2EpatternMatches_2EPMATCH\ A_27a\ A_27b \in ((A_27a^{(ty_2Elist_2Elist\ ((ty_2Eoption_2Eoption\ A_27a)^{A_27b})})})^{A_27b}) \quad (5)$$

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o\ (p\ P \Rightarrow p\ Q)$ of type ι .

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c_2Ebool_2E_21\ 2)\ (\lambda V2t \in 2.$

Definition 10 We define $c_2EpatternMatches_2EPMATCH_EQUIV_ROWS$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0v \in A_27a.\lambda V1rows1 \in (ty_2Elist_2Elist\ ((ty_2Eoption_2Eoption\ A_27b)^{A_27a}))$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0x \in A_27a.((V0x = V0x) \Leftrightarrow True)) \quad (9)$$

Theorem 1

$$\forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\forall V0v \in A_27a.(\forall V1rows \in (ty_2Elist_2Elist\ ((ty_2Eoption_2Eoption\ A_27b)^{A_27a})).(p\ (ap\ (ap\ (ap\ (ap\ (c_2EpatternMatches_2EPMATCH_EQUIV_ROWS\ A_27a\ A_27b)\ V0v)\ V1rows)\ V1rows))))$$