

thm\_2EpatternMatches\_2EPMATCH\_\_EQUIV\_\_ROWS\_\_is\_\_equiv\_\_  
 (TMZQSGCMgaEFDVsxBYaKCmU-  
 cLPy9V8CfKh5)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A-27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Eoption\_2Eoption A0) \quad (1)$$

Let  $c\_2Eoption\_2EIS\_SOME : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.\lambda a.nonempty A.\lambda a \Rightarrow c\_2Eoption\_2EIS\_SOME A.\lambda a \in (2^{(ty\_2Eoption\_2Eoption A.\lambda a)}) \quad (2)$$

Let  $ty\_2Elist\_2Elist : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow nonempty (ty\_2Elist\_2Elist A0) \quad (3)$$

Let  $c\_2Elist\_2ELIST\_TO\_SET : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A.\lambda a.nonempty A.\lambda a \Rightarrow c\_2Elist\_2ELIST\_TO\_SET A.\lambda a \in ((2^{A.\lambda a})^{(ty\_2Elist\_2Elist A.\lambda a)}) \quad (4)$$

**Definition 7** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a. (\lambda V1f \in (2^{A\_27a}). (ap\ V1f\ V0x)))$

**Definition 8** We define  $c\_2Emin\_2E40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if}\ (\exists x \in A. p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x. x \in A \wedge p\ x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 9** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E40\ A)\ P)))$

Let  $c\_2EpatternMatches\_2EPMATCH : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a. nonempty\ A\_27a \Rightarrow \forall A\_27b. nonempty\ A\_27b \Rightarrow c\_2EpatternMatches\_2EPMATCH \\ A\_27a\ A\_27b \in ((A\_27a^{(ty\_2Elist\_2Elist\ ((ty\_2Eoption\_2Eoption\ A\_27a)^{A\_27b})})})^{A\_27b} \end{aligned} \quad (5)$$

**Definition 10** We define  $c\_2Ebool\_2E2F\_5C$  to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap\ (c\_2Ebool\_2E21\ 2)\ (\lambda V2t \in 2. (ap\ (c\_2Ebool\_2E21\ 2)\ V2t)\ V1t2))))$

**Definition 11** We define  $c\_2EpatternMatches\_2EPMATCH\_EQUIV\_ROWS$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0v \in A\_27a. \lambda V1rows1 \in (ty\_2Elist\_2Elist\ ((ty\_2Eoption\_2Eoption\ A\_27b)^{A\_27a}))$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t) \Leftrightarrow (p\ V1x))) \Leftrightarrow (p\ V0t))) \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2. (((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow \\ True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (( \\ (p\ V0t) \Rightarrow False) \Leftrightarrow (\neg (p\ V0t)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (10)$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (11)$$

Assume the following.

$$\begin{aligned} (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow \\ ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \end{aligned} \quad (12)$$

Assume the following.

$$2.(((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))) \Rightarrow \quad (13)$$

$$(((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27))))$$

**Theorem 1**

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow \forall A\_27b.nonempty \ A\_27b \Rightarrow ($$

$$\forall V0v \in A\_27a. (\forall V1rows1 \in (ty\_2Elist\_2Elist ((ty\_2Eoption\_2Eoption$$

$$A\_27b)^{A\_27a})). (\forall V2rows2 \in (ty\_2Elist\_2Elist ((ty\_2Eoption\_2Eoption$$

$$A\_27b)^{A\_27a})). (\forall V3rows3 \in (ty\_2Elist\_2Elist ((ty\_2Eoption\_2Eoption$$

$$A\_27b)^{A\_27a})). ((p \ (ap \ (ap \ (ap \ (c\_2EpatternMatches\_2EPMATCH\_EQUIV\_ROWS$$

$$A\_27a \ A\_27b) \ V0v) \ V1rows1) \ V2rows2)) \Rightarrow ((p \ (ap \ (ap \ (ap \ (c\_2EpatternMatches\_2EPMATCH\_EQUIV\_ROWS$$

$$A\_27a \ A\_27b) \ V0v) \ V2rows2) \ V3rows3)) \Rightarrow (p \ (ap \ (ap \ (ap \ (c\_2EpatternMatches\_2EPMATCH\_EQUIV\_ROWS$$

$$A\_27a \ A\_27b) \ V0v) \ V1rows1) \ V3rows3))))))$$