

thm\_2EpatternMatches\_2EPMATCH\_ROW\_REDUNDANT\_NIL  
(TMWv7eD71R7LkZmJ1KUEXTfixDNsdfEVwCY)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 5** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num$

Let  $ty\_2Elist\_2Elist : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Elist\_2Elist\ A0) \tag{5}$$

Let  $c\_2Elist\_2ECONS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Elist\_2ECONS\ A\_27a \in (((ty\_2Elist\_2Elist\ A\_27a)^{(ty\_2Elist\_2Elist\ A\_27a)})^{A\_27a}) \tag{6}$$

Let  $c\_2Elist\_2ENIL : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Elist\_2ENIL\ A\_27a \in (ty\_2Elist\_2Elist\ A\_27a) \quad (7)$$

**Definition 6** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A$ .if  $(\exists x \in A.p\ (ap\ P\ x))$  then (the  $(\lambda x.x \in A \wedge p\ x)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 7** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40\ A\_27a\ V0P))))$

Let  $c\_2Elist\_2EEL : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Elist\_2EEL\ A\_27a \in ((A\_27a^{(ty\_2Elist\_2Elist\ A\_27a)})^{ty\_2Enum\_2Enum}) \quad (8)$$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Eoption\_2Eoption\ A0) \quad (9)$$

Let  $c\_2Eoption\_2EIS\_SOME : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2EIS\_SOME\ A\_27a \in (2^{(ty\_2Eoption\_2Eoption\ A\_27a)}) \quad (10)$$

**Definition 8** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o\ (p\ P \Rightarrow p\ Q)$  of type  $\iota$ .

Let  $c\_2Elist\_2ELENGTH : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Elist\_2ELENGTH\ A\_27a \in (ty\_2Enum\_2Enum^{(ty\_2Elist\_2Elist\ A\_27a)}) \quad (11)$$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in 2.(ap\ (c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ V2t))))$

**Definition 10** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_21\ 2))$

**Definition 11** We define  $c\_2Eprim\_rec\_2E\_3C$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.\lambda V1n \in ty\_2Enum\_2Enum$

**Definition 12** We define  $c\_2EpatternMatches\_2EPMATCH\_ROW\_REDUNDANT$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0v \in A\_27a.\lambda V1rs \in (ty\_2Elist\_2Elist\ ((ty\_2Eoption\_2Eoption\ A\_27b)^{A\_27a}))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \quad (12)$$

**Definition 13** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Assume the following.

$$True \quad (13)$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (14)$$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p V0t))) \quad (15)$$

Assume the following.

$$(\forall V0t \in 2. (((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \quad (16)$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \quad (17)$$

Assume the following.

$$\forall A\_27a. nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (18)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (19)$$

Assume the following.

$$\forall A\_27a. nonempty A\_27a \Rightarrow (((ap (c\_2Elist\_2ELENGTH A\_27a) (c\_2Elist\_2ENIL A\_27a)) = c\_2Enum\_2E0) \wedge (\forall V0h \in A\_27a. (\forall V1t \in (ty\_2Elist\_2Elist A\_27a). ((ap (c\_2Elist\_2ELENGTH A\_27a) (ap (ap (c\_2Elist\_2ECONS A\_27a) V0h) V1t)) = (ap c\_2Enum\_2ESUC (ap (c\_2Elist\_2ELENGTH A\_27a) V1t)))))) \quad (20)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum. (\neg(p (ap (ap c\_2Eprim\_rec\_2E\_3C V0n) c\_2Enum\_2E0)))) \quad (21)$$

### Theorem 1

$$\forall A\_27a. nonempty A\_27a \Rightarrow \forall A\_27b. nonempty A\_27b \Rightarrow (\forall V0v \in A\_27a. (\forall V1i \in ty\_2Enum\_2Enum. ((p (ap (ap (ap (c\_2EpatternMatches\_2EPMATCH\_ROW\_REDUNDANT A\_27a A\_27b) V0v) (c\_2Elist\_2ENIL ((ty\_2Eoption\_2Eoption A\_27b)^{A\_27a})) V1i)) \Leftrightarrow False)))$$