

# thm\_2EpatternMatches\_2Esome\_\_var\_\_bool\_\_F (TMcd6sStfxAUW6GvVoStYBVt6ks6taZR8hG)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_2E$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_2E1$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_2EF$  to be  $(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_2E7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_2EF$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \tag{1}$$

**Definition 7** We define  $c\_2Ebool\_2E\_2E5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_2E1 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \tag{2}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{3}$$

**Definition 8** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_2Esum\_2EABS\_sum$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Eoption\_2Eoption\ A0) \quad (4)$$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2Eoption\_ABS\ A\_27a \in ((ty\_2Eoption\_2Eoption\ A\_27a)^{(ty\_2Esum\_2Esum\ A\_27a\ ty\_2Eone\_2Eone)}) \quad (5)$$

**Definition 9** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A\_27a : \iota. \lambda V0x \in A\_27a. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ V0x)$

**Definition 10** We define  $c\_2Emin\_2E40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if}\ (\exists x \in A. P\ x)\ \mathbf{then}\ (the\ (\lambda x. x \in A)\ P)$  of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Eone\_2Eone$  to be  $(ap\ (c\_2Emin\_2E40\ ty\_2Eone\_2Eone)\ (\lambda V0x \in ty\_2Eone\_2Eone. V0x))$

**Definition 12** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27b. (ap\ (c\_2Esum\_2EABS\ A\_27a\ A\_27b)\ V0e)$

**Definition 13** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A\_27a : \iota. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ (c\_2Eone\_2Eone\ A\_27a))$

**Definition 14** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E40\ A\_27a)\ V0P)))$

**Definition 15** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (ap\ V2t2\ (ap\ V1t1\ V2t2))))))$

**Definition 16** We define  $c\_2Eoption\_2ESome$  to be  $\lambda A\_27a : \iota. \lambda V0P \in (2^{A\_27a}). (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27a)\ V0P)\ (c\_2Eoption\_2ENONE\ A\_27a))\ (c\_2Eoption\_2ESOME\ A\_27a))$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \quad (8)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0y \in A\_27a. (((ap\ (c\_2Eoption\_2ESome\ A\_27a)\ (\lambda V1x \in A\_27a. (ap\ (ap\ (c\_2Emin\_2E3D\ A\_27a)\ V1x)\ V0y))) = \\ (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0y)) \wedge ((ap\ (c\_2Eoption\_2ESome\ A\_27a)\ (\lambda V2x \in A\_27a. (ap\ (ap\ (c\_2Emin\_2E3D\ A\_27a)\ V2x)\ V0y))) = \\ (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0y)))))) \quad (9) \end{aligned}$$

**Theorem 1**

$$((ap\ (c\_2Eoption\_2ESome\ 2)\ (\lambda V0x \in 2. (ap\ c\_2Ebool\_2E7E\ V0x))) = (ap\ (c\_2Eoption\_2ESOME\ 2)\ c\_2Ebool\_2E7E))$$