

# thm\_2EpatternMatches\_2Esome\_\_var\_\_bool\_\_T (TMNfsczei6Z87HaDjAhm5KPZiQQJqbMjUGo)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2E\_EF$

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \tag{1}$$

**Definition 7** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Esum\_2Esum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow \forall A1.nonempty\ A1 \Rightarrow nonempty\ (ty\_2Esum\_2Esum\ A0\ A1) \tag{2}$$

Let  $c\_2Esum\_2EABS\_sum : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EABS\_sum\ A\_27a\ A\_27b \in ((ty\_2Esum\_2Esum\ A\_27a\ A\_27b)^{((2^{A\_27b})^{A\_27a})^2}) \tag{3}$$

**Definition 8** We define  $c\_2Esum\_2EINL$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27a.(ap (c\_2Esum\_2EABS\_sum$

Let  $ty\_2Eoption\_2Eoption : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty\ A0 \Rightarrow nonempty\ (ty\_2Eoption\_2Eoption\ A0) \quad (4)$$

Let  $c\_2Eoption\_2Eoption\_ABS : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow c\_2Eoption\_2Eoption\_ABS\ A\_27a \in ((ty\_2Eoption\_2Eoption\ A\_27a)^{(ty\_2Esum\_2Esum\ A\_27a\ ty\_2Eone\_2Eone)}) \quad (5)$$

**Definition 9** We define  $c\_2Eoption\_2ESOME$  to be  $\lambda A\_27a : \iota. \lambda V0x \in A\_27a. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ V0x)$

**Definition 10** We define  $c\_2Emin\_2E40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if}\ (\exists x \in A. P\ x)\ \mathbf{then}\ (the\ (\lambda x. x \in A)\ P)$  of type  $\iota \Rightarrow \iota$ .

**Definition 11** We define  $c\_2Eone\_2Eone$  to be  $(ap\ (c\_2Emin\_2E40\ ty\_2Eone\_2Eone)\ (\lambda V0x \in ty\_2Eone\_2Eone. V0x))$

**Definition 12** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0e \in A\_27b. (ap\ (c\_2Esum\_2EABS\ A\_27a)\ V0e)$

**Definition 13** We define  $c\_2Eoption\_2ENONE$  to be  $\lambda A\_27a : \iota. (ap\ (c\_2Eoption\_2Eoption\_ABS\ A\_27a)\ 0)$

**Definition 14** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota. (\lambda V0P \in (2^{A\_27a}). (ap\ V0P\ (ap\ (c\_2Emin\_2E40\ A\_27a)\ V0P)))$

**Definition 15** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. (ap\ V2t2\ (ap\ (c\_2Emin\_2E40\ A\_27a)\ V1t1)))))$

**Definition 16** We define  $c\_2Eoption\_2ESome$  to be  $\lambda A\_27a : \iota. \lambda V0P \in (2^{A\_27a}). (ap\ (ap\ (ap\ (c\_2Ebool\_2ECOND\ A\_27a)\ V0P)\ c\_2Eoption\_2ENONE))$

Assume the following.

$$True \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow \neg(p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow \neg(p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0y \in A\_27a. (((ap\ (c\_2Eoption\_2ESome\ A\_27a)\ (\lambda V1x \in A\_27a. (ap\ (ap\ (c\_2Emin\_2E3D\ A\_27a)\ V1x)\ V0y))) = \\ & (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0y)) \wedge ((ap\ (c\_2Eoption\_2ESome\ A\_27a)\ (\lambda V2x \in A\_27a. (ap\ (ap\ (c\_2Emin\_2E3D\ A\_27a)\ V2x)\ V0y))) = \\ & (ap\ (c\_2Eoption\_2ESOME\ A\_27a)\ V0y)))) \end{aligned} \quad (9)$$

**Theorem 1**

$$((ap\ (c\_2Eoption\_2ESome\ 2)\ (\lambda V0x \in 2. V0x)) = (ap\ (c\_2Eoption\_2ESOME\ 2)\ c\_2Ebool\_2ET))$$