

thm_2Epred_set_2EEMPTY_SUBSET
(TMcv283nGah117GZ3Zn6t79HqjhNJyCmBA4)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E21 2) (\lambda V2t \in 2.V2t))$

Definition 7 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2EF)$.

Definition 8 We define $c_2Ebool_2E_2E_27E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2EF$

Definition 9 We define $c_2Ebool_2E_2EIN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 10 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_27a.(p V0t) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$\begin{aligned}
& (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\
& True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\
& (p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t))))))
\end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& \forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.(\neg(p (ap (ap \\
& (c.2Ebool_2EIN A.27a) V0x) (c.2Epred_set.2EEMPTY A.27a))))))
\end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned}
& \forall A.27a.nonempty A.27a \Rightarrow (\forall V0s \in (2^{A.27a}).(p (ap (\\
& ap (c.2Epred_set.2ESUBSET A.27a) (c.2Epred_set.2EEMPTY A.27a)) \\
& V0s)))
\end{aligned}$$