

thm_2Epred_set_INSERT_applied (TMRrpULgJ2AwEvJV4nWa24ftfhEuphJLZQx)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 4 We define `c_2Emarker_2E_Cong` to be $\lambda V0x \in 2. V0x$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))))$

Definition 7 We define `c_2Ebool_2E_IN` to be $\lambda A. 27a : \iota. (\lambda V0x \in A. 27a. (\lambda V1f \in (2^{A-27a}). (\text{ap } V1f \ V0x))))$

Definition 8 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))))$

Let `ty_2Epair_2Eprod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (ty_2Epair_2Eprod \ A0 \ A1) \tag{1}$$

Let `c_2Epair_2EABS_prod` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow \forall A. 27b. \text{nonempty } A. 27b \Rightarrow c_2Epair_2EABS_prod \ A. 27a \ A. 27b \in ((ty_2Epair_2Eprod \ A. 27a \ A. 27b)^{(2^{A-27b})^{A-27a}}) \tag{2}$$

Definition 9 We define `c_2Epair_2E_2C` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0x \in A. 27a. \lambda V1y \in A. 27b. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2z \in 2. V2z))$

Let `c_2Epred_set_2EGSPEC` : $\iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow \forall A. 27b. \text{nonempty } A. 27b \Rightarrow c_2Epred_set_2EGSPEC \ A. 27a \ A. 27b \in ((2^{A-27a})^{(ty_2Epair_2Eprod \ A. 27a \ 2)^{A-27b}}) \tag{3}$$

Definition 10 We define $c_2Epred_set_2EINSERT$ to be $\lambda A_27a : \iota.\lambda V0x \in A_27a.\lambda V1s \in (2^{A_27a}).(ap (c_2Ebool_2EIN A_27a) (p (ap (ap (c_2Ebool_2EIN A_27a) V0x) (ap (ap (c_2Epred_set_2EINSERT A_27a) V1y) V2s)))) \Leftrightarrow ((V0x = V1y) \vee (p (ap (ap (c_2Ebool_2EIN A_27a) V0x) V2s))))))$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in \\ & A_27a. (\forall V2s \in (2^{A_27a}). ((p (ap (ap (c_2Ebool_2EIN A_27a) \\ & V0x) (ap (ap (c_2Epred_set_2EINSERT A_27a) V1y) V2s)))) \Leftrightarrow ((V0x = \\ & V1y) \vee (p (ap (ap (c_2Ebool_2EIN A_27a) V0x) V2s))))))) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a. (\forall V1y \in \\ & A_27a. (\forall V2s \in (2^{A_27a}). ((p (ap (ap (ap (c_2Epred_set_2EINSERT \\ & A_27a) V1y) V2s) V0x)) \Leftrightarrow ((V0x = V1y) \vee (p (ap (ap (c_2Ebool_2EIN A_27a) \\ & V0x) V2s))))))) \end{aligned}$$