

# thm\_2Epred\_set\_2EINTER\_COMM (TMJrQkT- NyvrtFbQ4T1KRdZMtSwE4uiAKnQS)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2E\_2T$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 5** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2$

**Definition 6** We define  $c\_2Ebool\_2E\_2IN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \tag{1}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{2}$$

**Definition 7** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2E$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC A\_27a A\_27b \in ((2^{A\_27a})^{((ty\_2Epair\_2Eprod A\_27a 2)^{A\_27b})}) \tag{3}$$

**Definition 8** We define  $c\_2Epred\_set\_2EINTER$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap (c\_2$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \wedge (p V1t2)) \Leftrightarrow ((p V1t2) \wedge (p V0t1)))))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty A.27a \Rightarrow (\forall V0s \in (2^{A.27a}).(\forall V1t \in \\ & (2^{A.27a}).((V0s = V1t) \Leftrightarrow (\forall V2x \in A.27a.((p (ap (ap (c.2Ebool.2EIN \\ & A.27a) V2x) V0s)) \Leftrightarrow (p (ap (ap (c.2Ebool.2EIN A.27a) V2x) V1t))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall A.27a.nonempty A.27a \Rightarrow (\forall V0s \in (2^{A.27a}).(\forall V1t \in \\ & (2^{A.27a}).(\forall V2x \in A.27a.((p (ap (ap (c.2Ebool.2EIN A.27a) \\ & V2x) (ap (ap (c.2Epred\_set.2EINTER A.27a) V0s) V1t))) \Leftrightarrow ((p (ap \\ & (ap (c.2Ebool.2EIN A.27a) V2x) V0s)) \wedge (p (ap (ap (c.2Ebool.2EIN \\ & A.27a) V2x) V1t))))))) \end{aligned} \quad (6)$$

**Theorem 1**

$$\begin{aligned} & \forall A.27a.nonempty A.27a \Rightarrow (\forall V0s \in (2^{A.27a}).(\forall V1t \in \\ & (2^{A.27a}).((ap (ap (c.2Epred\_set.2EINTER A.27a) V0s) V1t) = ( \\ & ap (ap (c.2Epred\_set.2EINTER A.27a) V1t) V0s)))) \end{aligned}$$