

# thm\_2Epred\_set\_2EMEMBER\_NOT\_EMPTY (TMVJKKQ9rybekc5Jit5FWj1eA3oGKdxYL81)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_40` to be  $\lambda A.\lambda P \in 2^A.$ **if**  $(\exists x \in A.p (ap P x))$  **then** (the  $(\lambda x.x \in A \wedge p x)$  of type  $\iota \Rightarrow \iota$ ).

**Definition 2** We define `c_2Emin_2E_3D` to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define `c_2Ebool_2E_3F` to be  $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap V0P (ap (c_2Emin_2E_40 A_{27a}) P)))$

**Definition 4** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 5** We define `c_2Ebool_2E_T` to be  $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 6** We define `c_2Ebool_2E_21` to be  $\lambda A_{27a} : \iota.(\lambda V0P \in (2^{A_{27a}}).(ap (ap (c_2Emin_2E_3D (2^{A_{27a}})) P)))$

**Definition 7** We define `c_2Ebool_2E_F` to be  $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 8** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2.V2t)))$

**Definition 9** We define `c_2Epred_set_2EEMPTY` to be  $\lambda A_{27a} : \iota.(\lambda V0x \in A_{27a}.c_2Ebool_2E_F)$ .

**Definition 10** We define `c_2Ebool_2E_IN` to be  $\lambda A_{27a} : \iota.(\lambda V0x \in A_{27a}.(\lambda V1f \in (2^{A_{27a}}).(ap V1f V0x)))$

**Definition 11** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_21))$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (4)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (5)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t))))) \quad (7)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0s \in (2^{A\_27a}). (\forall V1t \in (2^{A\_27a}). ((V0s = V1t) \Leftrightarrow (\forall V2x \in A\_27a. ((p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A\_27a)\ V2x)\ V0s)) \Leftrightarrow (p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A\_27a)\ V2x)\ V1t)))))) \quad (8)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\neg(p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A\_27a)\ V0x)\ (c\_2Epred\_set\_2EEMPTY\ A\_27a)))) \quad (9)$$

**Theorem 1**

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0s \in (2^{A\_27a}). ((\exists V1x \in A\_27a. (p\ (ap\ (ap\ (c\_2Ebool\_2EIN\ A\_27a)\ V1x)\ V0s))) \Leftrightarrow (\neg(V0s = (c\_2Epred\_set\_2EEMPTY\ A\_27a))))$$