

# thm\_2Epred\_set\_2ENOT\_PSUBSET\_EMPTY (TMGivqeZve9x5Mvw8KBR4rStFjb6nLc5NSK)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2EF)$ .

**Definition 7** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 8** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 9** We define  $c\_2Epred\_set\_2ESUBSET$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap (c$

**Definition 10** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in$

**Definition 11** We define  $c\_2Epred\_set\_2EPSUBSET$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).\lambda V1t \in (2^{A\_27a}).(ap$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2. (\neg((p\ V0t) \wedge (\neg(p\ V0t)))))) \quad (5)$$

Assume the following.

$$((\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0s \in (2^{A\_27a}). ((p\ (ap\ (ap\ (c\_2Epred\_set\_2ESUBSET\ A\_27a)\ V0s)\ (c\_2Epred\_set\_2EEMPTY\ A\_27a))) \Leftrightarrow (V0s = (c\_2Epred\_set\_2EEMPTY\ A\_27a)))) \quad (7)$$

**Theorem 1**

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0s \in (2^{A\_27a}). (\neg(p\ (ap\ (ap\ (c\_2Epred\_set\_2EPSUBSET\ A\_27a)\ V0s)\ (c\_2Epred\_set\_2EEMPTY\ A\_27a))))))$$