

thm_2Epred_set_2ESUBSET__ANTISYM (TMY- Jav3SeFZNQ6kwVrJnQYKJhxZ4qhMbmmdJ)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 5 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 6 We define $c_2Ebool_2E_2IN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x)))$

Definition 7 We define $c_2Epred_set_2ESUBSET$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).\lambda V1t \in (2^{A_27a}).(ap (c$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \quad (1)$$

Assume the following.

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0s \in (2^{A_27a}).(\forall V1t \in (2^{A_27a}).((V0s = V1t) \Leftrightarrow (\forall V2x \in A_27a.((p (ap (ap (c_2Ebool_2E_2IN A_27a) V2x) V0s)) \Leftrightarrow (p (ap (ap (c_2Ebool_2E_2IN A_27a) V2x) V1t))))))) \quad (2)$$

Theorem 1

$$\forall A_27a.nonempty A_27a \Rightarrow (\forall V0s \in (2^{A_27a}).(\forall V1t \in (2^{A_27a}).(((p (ap (ap (c_2Epred_set_2ESUBSET A_27a) V0s) V1t)) \wedge (p (ap (ap (c_2Epred_set_2ESUBSET A_27a) V1t) V0s))) \Rightarrow (V0s = V1t))))$$