

thm_2Epred_set_2ESUBSET_REFL (TM- LAALVsj2JepPGDXKAnrTWRkfsbKr33SmA)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. \lambda 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a})) (\lambda V1P \in 2. V1P)) (\lambda V2P \in 2. V2P)))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2. V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) (\text{c_2Ebool_2E_2F } V0t)) (\lambda V1t \in 2. V1t)))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))) (\lambda V3t \in 2. V3t)))$

Definition 8 We define `c_2Ebool_2EIN` to be $\lambda A. \lambda 27a : \iota. (\lambda V0x \in A. \lambda 27a. (\lambda V1f \in (2^{A-27a}). (\text{ap } V1f V0x))) (\lambda V2f \in (2^{A-27a}). (\text{ap } V2f V0x)))$

Definition 9 We define `c_2Epred_set_2ESUBSET` to be $\lambda A. \lambda 27a : \iota. \lambda V0s \in (2^{A-27a}). \lambda V1t \in (2^{A-27a}). (\text{ap } (\text{c_2Ebool_2EIN } A. 27a) V0s) V1t)$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$\forall A. \lambda 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A. 27a. (p V0t) \Leftrightarrow (p V1x))) \Leftrightarrow (p V0t))) \tag{2}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((\text{True} \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow \text{True}) \Leftrightarrow \\ & \text{True}) \wedge (((\text{False} \Rightarrow (p V0t)) \Leftrightarrow \text{True}) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow \text{True}) \wedge ((\\ & (p V0t) \Rightarrow \text{False}) \Leftrightarrow (\neg (p V0t)))))) \end{aligned} \tag{3}$$

Theorem 1

$$\forall A. \lambda 27a. \text{nonempty } A. 27a \Rightarrow (\forall V0s \in (2^{A-27a}). (p (\text{ap } (\text{c_2Epred_set_2ESUBSET } A. 27a) V0s) V0s)))$$