

thm_2Epred__set_2Ecountable__EMPTY
(TMQkEAGWVpZCbLCzWTjhBpkCjiwVY-
GoLFS6)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2ET$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}))$

Definition 4 We define $c_2Ebool_2E_2EF$ to be $(ap (c_2Ebool_2E_2E21 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_2E7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2EF$

Definition 7 We define $c_2Epred_set_2EEMPTY$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2EF)$.

Definition 8 We define $c_2Ebool_2E_2E2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E21 2) (\lambda V2t \in 2.V2t))$

Definition 9 We define $c_2Epred_set_2EUNIV$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.c_2Ebool_2E_2ET)$.

Definition 10 We define $c_2Ebool_2E_2EIN$ to be $\lambda A_27a : \iota.(\lambda V0x \in A_27a.(\lambda V1f \in (2^{A_27a}).(ap V1f V0x))$

Definition 11 We define $c_2Epred_set_2EINJ$ to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).\lambda V1s \in (2^{A_27b})$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Definition 12 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 13 We define $c_2Ebool_2E_2E3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40$

Definition 14 We define $c_2Epred_set_2Ecountable$ to be $\lambda A_27a : \iota.\lambda V0s \in (2^{A_27a}).(ap (c_2Ebool_2E_2E3F$

Assume the following.

$$True \quad (2)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A_{.27a}.(p\ V0t)) \Leftrightarrow (p\ V0t))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (4)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow \forall A_{.27b}.nonempty\ A_{.27b} \Rightarrow (\forall V0f \in (A_{.27b}^{A_{.27a}}). ((\forall V1s \in (2^{A_{.27b}}). (p\ (ap\ (ap\ (ap\ (c_{.2Epred_set_2EINJ}\ A_{.27a}\ A_{.27b})\ V0f)\ (c_{.2Epred_set_2EEMPTY}\ A_{.27a}))\ V1s))) \wedge (\forall V2s \in (2^{A_{.27a}}). ((p\ (ap\ (ap\ (ap\ (c_{.2Epred_set_2EINJ}\ A_{.27a}\ A_{.27b})\ V0f)\ V2s)\ (c_{.2Epred_set_2EEMPTY}\ A_{.27b}))) \Leftrightarrow (V2s = (c_{.2Epred_set_2EEMPTY}\ A_{.27a})))))) \quad (5)$$

Theorem 1

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (p\ (ap\ (c_{.2Epred_set_2Ecountable}\ A_{.27a})\ (c_{.2Epred_set_2EEMPTY}\ A_{.27a})))$$