

thm\_2Epred\_\_set\_2Enum\_\_countable  
 (TMGUXyemsKw-  
 pHmFhws4RKB4VaP9VY3TzY1y)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 9** We define  $c\_2Epred\_set\_2EUNIV$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.c\_2Ebool\_2ET)$ .

**Definition 10** We define  $c\_2Epred\_set\_2EINJ$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0f \in (A\_27b^{A\_27a}).\lambda V1s \in (2^{A\_27b})$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

**Definition 11** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \text{ then } (the (\lambda x.x \in A \wedge p (ap P x)))$  of type  $\iota \Rightarrow \iota$ .

**Definition 12** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E\_40$

**Definition 13** We define  $c\_2Epred\_set\_2Ecountable$  to be  $\lambda A\_27a : \iota.\lambda V0s \in (2^{A\_27a}).(ap (c\_2Ebool\_2E\_3F$

Assume the following.

$$True \quad (2)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a.(p \ V0t)) \Leftrightarrow (p \ V0t))) \quad (3)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge True) \Leftrightarrow \\ & (p \ V0t)) \wedge (((False \wedge (p \ V0t)) \Leftrightarrow False) \wedge (((p \ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Rightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (((p \ V0t) \Rightarrow (p \ V0t)) \Leftrightarrow True) \wedge (( \\ & (p \ V0t) \Rightarrow False) \Leftrightarrow (\neg(p \ V0t)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a. ((V0x = V0x) \Leftrightarrow True)) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a. (\forall V1y \in A\_27a. ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (7)$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p \ V0t)) \wedge (((False \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg(p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow False) \Leftrightarrow (\neg \\ & (p \ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p \ V0t1) \Rightarrow \\ & ((p \ V1t2) \Rightarrow (p \ V2t3))) \Leftrightarrow (((p \ V0t1) \wedge (p \ V1t2)) \Rightarrow (p \ V2t3)))))) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x\_27 \in 2. (\forall V2y \in 2. (\forall V3y\_27 \in \\ & 2. (((p \ V0x) \Leftrightarrow (p \ V1x\_27)) \wedge ((p \ V1x\_27) \Rightarrow ((p \ V2y) \Leftrightarrow (p \ V3y\_27)))) \Rightarrow \\ & (((p \ V0x) \Rightarrow (p \ V2y)) \Leftrightarrow ((p \ V1x\_27) \Rightarrow (p \ V3y\_27)))))) \end{aligned} \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty \ A\_27a \Rightarrow (\forall V0x \in A\_27a. (p \ (ap \ (ap \ (c\_2Ebool\_2EIN \ A\_27a) \ V0x) \ (c\_2Epred\_set\_2EUNIV \ A\_27a)))) \quad (11)$$

**Theorem 1**

$$(p \ (ap \ (c\_2Epred\_set\_2Ecountable \ ty\_2Enum\_2Enum) \ (c\_2Epred\_set\_2EUNIV \ ty\_2Enum\_2Enum)))$$