

thm_2Eprim_rec_2ELESS_NOT_EQ (TMVZn3Du55FzC51GysjHywAj57U5KC1Ds9J)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 2 We define $c_2Ebool_2E_2E$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define $c_2Ebool_2E_2E$ to be $\lambda A.\lambda a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) (\lambda V1P \in 2.V1P)) (\lambda V2P \in 2.V2P))$

Definition 4 We define $c_2Ebool_2E_2E$ to be $(ap (c_2Ebool_2E_2E 2) (\lambda V0t \in 2.V0t))$.

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 6 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E_2E))$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_2E 2) (\lambda V2t \in 2.V2t))$

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (ap c_2Enum_2EREP_num (ap c_2Enum_2ESUC_REP m)))$

Definition 9 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x)) \mathbf{then} (the (\lambda x.x \in A \wedge P x))$ of type $\iota \Rightarrow \iota$.

Definition 10 We define `c_2Ebool_2E_3F` to be $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (ap\ V0P\ (ap\ (c_2Emin_2E_40$

Definition 11 We define `c_2Eprim_rec_2E_3C` to be $\lambda V0m \in ty_2Enum_2Enum. \lambda V1n \in ty_2Enum_2Enum$

Assume the following.

$$(\forall V0t \in 2. (False \Rightarrow (p\ V0t))) \tag{5}$$

Assume the following.

$$(\forall V0n \in ty_2Enum_2Enum. (\neg (p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V0n)\ V0n)))) \tag{6}$$

Theorem 1

$$(\forall V0m \in ty_2Enum_2Enum. (\forall V1n \in ty_2Enum_2Enum. (p\ (ap\ (ap\ c_2Eprim_rec_2E_3C\ V0m)\ V1n)) \Rightarrow (\neg (V0m = V1n))))$$