

thm\_2Eprim\_rec\_2ETC\_\_IM\_\_RTC\_\_SUC  
(TMK1qhZDWQNZj1DM6PFjMaSAioyCKJKfnmX)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow p \Rightarrow Q)$  of type  $\iota$ .

**Definition 2** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 3** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$ .

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a})))$

**Definition 5** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{1}$$

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{2}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{3}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{4}$$

**Definition 7** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap c\_2Enum\_2EABS\_num$

**Definition 8** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 9** We define  $c\_2Erelation\_2ERTC$  to be  $\lambda A\_27a : \iota.\lambda V0R \in ((2^{A\_27a})^{A\_27a}).\lambda V1a \in A\_27a.\lambda V2b \in A\_27a.$

**Definition 10** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \ x)) \text{ then (the } (\lambda x. x \in A) \text{ of type } \iota \Rightarrow \iota.$

**Definition 11** We define `c_2Ebool_2E_3F` to be  $\lambda A_{.27a} : \iota. (\lambda V0P \in (2^{A_{.27a}}). (\text{ap } V0P \ (\text{ap } (\text{c}_2\text{Emin}_2\text{E}_40$

**Definition 12** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c}_2\text{Ebool}_2\text{E}_21 \ 2) \ (\lambda V2t \in$

**Definition 13** We define `c_2Erelation_2ETC` to be  $\lambda A_{.27a} : \iota. \lambda V0R \in ((2^{A_{.27a}})^{A_{.27a}}). \lambda V1a \in A_{.27a}. \lambda V2b$

Assume the following.

$$\text{True} \tag{5}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \ V0t1) \Rightarrow (p \ V1t2)) \Rightarrow (((p \ V1t2) \Rightarrow (p \ V0t1)) \Rightarrow ((p \ V0t1) \Leftrightarrow (p \ V1t2)))))) \tag{6}$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \wedge \text{True}) \Leftrightarrow (p \ V0t)) \wedge (((\text{False} \wedge (p \ V0t)) \Leftrightarrow \text{False}) \wedge (((p \ V0t) \wedge \text{False}) \Leftrightarrow \text{False}) \wedge (((p \ V0t) \wedge (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \vee (p \ V0t)) \Leftrightarrow \text{True}) \wedge (((p \ V0t) \vee \text{True}) \Leftrightarrow \text{True}) \wedge (((\text{False} \vee (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \vee \text{False}) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \vee (p \ V0t)) \Leftrightarrow (p \ V0t)))))) \tag{8}$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}. ((V0x = V0x) \Leftrightarrow \text{True})) \tag{9}$$

Assume the following.

$$(\forall V0t \in 2. (((\text{True} \Leftrightarrow (p \ V0t)) \Leftrightarrow (p \ V0t)) \wedge (((p \ V0t) \Leftrightarrow \text{True}) \Leftrightarrow (p \ V0t)) \wedge (((\text{False} \Leftrightarrow (p \ V0t)) \Leftrightarrow (\neg (p \ V0t))) \wedge (((p \ V0t) \Leftrightarrow \text{False}) \Leftrightarrow (\neg (p \ V0t)))))) \tag{10}$$

Assume the following.

$$(\forall V0m \in \text{ty}_2\text{Enum}_2\text{Enum}. (\forall V1n \in \text{ty}_2\text{Enum}_2\text{Enum}. ((\text{ap } \text{c}_2\text{Enum}_2\text{ESUC } V0m) = (\text{ap } \text{c}_2\text{Enum}_2\text{ESUC } V1n)) \Leftrightarrow (V0m = V1n)))) \tag{11}$$

Assume the following.

$$\forall A_{.27a}. \text{nonempty } A_{.27a} \Rightarrow (\forall V0R \in ((2^{A_{.27a}})^{A_{.27a}}). (\forall V1x \in A_{.27a}. (\forall V2y \in A_{.27a}. ((p \ (\text{ap } (\text{ap } (\text{ap } (\text{c}_2\text{Erelation}_2\text{ERTC } A_{.27a}) \ V0R) \ V1x) \ V2y)) \Leftrightarrow ((V1x = V2y) \vee (p \ (\text{ap } (\text{ap } (\text{ap } (\text{c}_2\text{Erelation}_2\text{ETC } A_{.27a}) \ V0R) \ V1x) \ V2y)))))) \tag{12}$$

Assume the following.

$$\begin{aligned}
& \forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0R \in ((2^{A_{27a}})^{A_{27a}}). \\
& (\forall V1x \in A_{27a}. (\forall V2z \in A_{27a}. ((p (ap (ap (ap (c\_2Erelation\_2ETC \\
& A_{27a}) V0R) V1x) V2z)) \Leftrightarrow ((p (ap (ap V0R V1x) V2z)) \vee (\exists V3y \in A_{27a}. \\
& ((p (ap (ap (ap (c\_2Erelation\_2ETC A_{27a}) V0R) V1x) V3y)) \wedge (p (ap \\
& (ap V0R V3y) V2z))))))))))
\end{aligned} \tag{13}$$

**Theorem 1**

$$\begin{aligned}
& (\forall V0m \in ty\_2Enum\_2Enum. (\forall V1n \in ty\_2Enum\_2Enum. ( \\
& (p (ap (ap (ap (c\_2Erelation\_2ETC ty\_2Enum\_2Enum) (\lambda V2x \in ty\_2Enum\_2Enum. \\
& (\lambda V3y \in ty\_2Enum\_2Enum. (ap (ap (c\_2Emin\_2E\_3D ty\_2Enum\_2Enum) \\
& V3y) (ap c\_2Enum\_2ESUC V2x)))))) V0m) (ap c\_2Enum\_2ESUC V1n)))) \Leftrightarrow \\
& (p (ap (ap (ap (c\_2Erelation\_2ERTC ty\_2Enum\_2Enum) (\lambda V4x \in ty\_2Enum\_2Enum. \\
& (\lambda V5y \in ty\_2Enum\_2Enum. (ap (ap (c\_2Emin\_2E\_3D ty\_2Enum\_2Enum) \\
& V5y) (ap c\_2Enum\_2ESUC V4x)))))) V0m) V1n))))
\end{aligned}$$