

thm\_2Eprim\_\_rec\_2EWF\_\_PRED  
(TMRA8Cbp5yDH7szDcCoKEo5sL5b7xPfG1M7)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E\_21$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E\_21$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t)))$

Let  $c\_2Enum\_2EZERO\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2EZERO\_REP \in \omega \tag{1}$$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{2}$$

Let  $c\_2Enum\_2EABS\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EABS\_num \in (ty\_2Enum\_2Enum^{\omega}) \tag{3}$$

**Definition 7** We define  $c\_2Enum\_2E0$  to be  $(ap\ c\_2Enum\_2EABS\_num\ c\_2Enum\_2EZERO\_REP)$ .

Let  $c\_2Enum\_2EREP\_num : \iota$  be given. Assume the following.

$$c\_2Enum\_2EREP\_num \in (\omega^{ty\_2Enum\_2Enum}) \tag{4}$$

Let  $c\_2Enum\_2ESUC\_REP : \iota$  be given. Assume the following.

$$c\_2Enum\_2ESUC\_REP \in (\omega^{\omega}) \tag{5}$$

**Definition 8** We define  $c\_2Enum\_2ESUC$  to be  $\lambda V0m \in ty\_2Enum\_2Enum.(ap\ c\_2Enum\_2EABS\_num$

**Definition 9** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap\ (ap\ c\_2Emin\_2E\_3D\_3D\_3E\ V0t)\ c\_2Ebool\_2E\_7E$

**Definition 10** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap\ (c\_2Ebool\_2E\_21\ 2)\ (\lambda V2t \in$

**Definition 11** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A.\lambda P \in 2^A.\mathbf{if}\ (\exists x \in A.p\ (ap\ P\ x))\ \mathbf{then}\ (the\ (\lambda x.x \in A \wedge$   
of type  $\iota \Rightarrow \iota$ .

**Definition 12** We define  $c\_2Ebool\_2E\_3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap\ V0P\ (ap\ (c\_2Emin\_2E\_40$

**Definition 13** We define  $c\_2Erelation\_2EWF$  to be  $\lambda A\_27a : \iota.\lambda V0R \in ((2^{A\_27a})^{A\_27a}).(ap\ (c\_2Ebool\_2E\_21$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \tag{7}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \tag{8}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \tag{9}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{10}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \tag{11}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).((\neg(\forall V1x \in A\_27a.(p\ (ap\ V0P\ V1x)))) \Leftrightarrow (\exists V2x \in A\_27a.(\neg(p\ (ap\ V0P\ V2x)))))) \tag{12}$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0P \in (2^{A\_27a}).((\neg(\exists V1x \in A\_27a.(p\ (ap\ V0P\ V1x)))) \Leftrightarrow (\forall V2x \in A\_27a.(\neg(p\ (ap\ V0P\ V2x)))))) \tag{13}$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.((\neg((p V0A) \Rightarrow (p V1B))) \Leftrightarrow ((p V0A) \wedge (\neg(p V1B))))) \quad (14)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \wedge (p V1B))) \Leftrightarrow ((\neg(p V0A) \vee \neg(p V1B)))) \wedge ((\neg((p V0A) \vee (p V1B))) \Leftrightarrow ((\neg(p V0A) \wedge \neg(p V1B))))) \quad (15)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((p V0A) \Rightarrow (p V1B)) \Leftrightarrow ((\neg(p V0A) \vee (p V1B)))) \quad (16)$$

Assume the following.

$$(\forall V0n \in ty\_2Enum\_2Enum.(\neg((ap\ c\_2Enum\_2ESUC\ V0n) = c\_2Enum\_2E0))) \quad (17)$$

Assume the following.

$$(\forall V0P \in (2^{ty\_2Enum\_2Enum}).(((p\ (ap\ V0P\ c\_2Enum\_2E0)) \wedge (\forall V1n \in ty\_2Enum\_2Enum.((p\ (ap\ V0P\ V1n)) \Rightarrow (p\ (ap\ V0P\ (ap\ c\_2Enum\_2ESUC\ V1n)))))) \Rightarrow (\forall V2n \in ty\_2Enum\_2Enum.(p\ (ap\ V0P\ V2n)))) \quad (18)$$

Assume the following.

$$(\forall V0m \in ty\_2Enum\_2Enum.(\forall V1n \in ty\_2Enum\_2Enum.((ap\ c\_2Enum\_2ESUC\ V0m) = (ap\ c\_2Enum\_2ESUC\ V1n)) \Leftrightarrow (V0m = V1n))) \quad (19)$$

**Theorem 1**

$$(p\ (ap\ (c\_2Erelation\_2EWF\ ty\_2Enum\_2Enum)\ (\lambda V0x \in ty\_2Enum\_2Enum. (\lambda V1y \in ty\_2Enum\_2Enum.(ap\ (ap\ (c\_2Emin\_2E\_3D\ ty\_2Enum\_2Enum)\ V1y)\ (ap\ c\_2Enum\_2ESUC\ V0x))))))$$