

thm_2Eprim__rec_2Enum__Axiom
(TMagsjrwFAi7CUQ7Qe5dQ4qytX3KNH24dNn)

October 26, 2020

Definition 1 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$ of type ι .

Definition 2 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p x)$) of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$ of type $\iota \Rightarrow \iota$.

Definition 4 We define $c_2Ebool_2E_3F$ to be $\lambda A.\lambda P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A P)))$

Definition 5 We define c_2Ebool_2ET to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A.\lambda P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27a})) P))$

Definition 7 We define c_2Ebool_2EF to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Let $ty_2Enum_2Enum : \iota$ be given. Assume the following.

$$nonempty\ ty_2Enum_2Enum \tag{1}$$

Let $c_2Enum_2EREP_num : \iota$ be given. Assume the following.

$$c_2Enum_2EREP_num \in (\omega^{ty_2Enum_2Enum}) \tag{2}$$

Let $c_2Enum_2ESUC_REP : \iota$ be given. Assume the following.

$$c_2Enum_2ESUC_REP \in (\omega^{\omega}) \tag{3}$$

Let $c_2Enum_2EABS_num : \iota$ be given. Assume the following.

$$c_2Enum_2EABS_num \in (ty_2Enum_2Enum^{\omega}) \tag{4}$$

Definition 8 We define c_2Enum_2ESUC to be $\lambda V0m \in ty_2Enum_2Enum.(ap c_2Enum_2EABS_num (c_2Enum_2ESUC_REP m))$

