

# thm\_2Eproduct\_2ENPRODUCT\_UNION (TMLQQurk6ZzY6eiJibhmnxfCxgahZ4nVML5)

October 26, 2020

**Definition 1** We define  $c\_2Emin\_2E\_3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2ET$  to be  $(ap (ap (c\_2Emin\_2E\_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E\_21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E\_3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2EF$  to be  $(ap (c\_2Ebool\_2E\_21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E\_3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E\_7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E\_3D\_3D\_3E V0t) c\_2Ebool\_2EF$

**Definition 7** We define  $c\_2Ebool\_2EIN$  to be  $\lambda A\_27a : \iota.(\lambda V0x \in A\_27a.(\lambda V1f \in (2^{A\_27a}).(ap V1f V0x)))$

**Definition 8** We define  $c\_2Ebool\_2E\_5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

**Definition 9** We define  $c\_2Ebool\_2E\_2F\_5C$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E\_21 2) (\lambda V2t \in 2.V2t))$

Let  $ty\_2Epair\_2Eprod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0.nonempty A0 \Rightarrow \forall A1.nonempty A1 \Rightarrow nonempty (ty\_2Epair\_2Eprod A0 A1) \tag{1}$$

Let  $c\_2Epair\_2EABS\_prod : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epair\_2EABS\_prod A\_27a A\_27b \in ((ty\_2Epair\_2Eprod A\_27a A\_27b)^{(2^{A\_27b})^{A\_27a}}) \tag{2}$$

**Definition 10** We define  $c\_2Epair\_2E\_2C$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0x \in A\_27a.\lambda V1y \in A\_27b.(ap (c\_2Ebool\_2EIN$

Let  $c\_2Epred\_set\_2EGSPEC : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow c\_2Epred\_set\_2EGSPEC A\_27a A\_27b \in ((2^{A\_27a})^{(ty\_2Epair\_2Eprod A\_27a 2)^{A\_27b}}) \tag{3}$$

**Definition 11** We define  $c\_2Epred\_set\_2EUNION$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). \lambda V1t \in (2^{A\_27a}). (ap (c\_2E$

**Definition 12** We define  $c\_2Epred\_set\_2EEMPTY$  to be  $\lambda A\_27a : \iota. (\lambda V0x \in A\_27a.c\_2Ebool\_2EF).$

**Definition 13** We define  $c\_2Epred\_set\_2EINTER$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). \lambda V1t \in (2^{A\_27a}). (ap (c\_2E$

**Definition 14** We define  $c\_2Epred\_set\_2EDISJOINT$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). \lambda V1t \in (2^{A\_27a}). (ap (c\_2E$

**Definition 15** We define  $c\_2Epred\_set\_2EINSERT$  to be  $\lambda A\_27a : \iota. \lambda V0x \in A\_27a. \lambda V1s \in (2^{A\_27a}). (ap (c\_2E$

**Definition 16** We define  $c\_2Epred\_set\_2EFINITE$  to be  $\lambda A\_27a : \iota. \lambda V0s \in (2^{A\_27a}). (ap (c\_2Ebool\_2E\_21 (2$

**Definition 17** We define  $c\_2Emin\_2E\_40$  to be  $\lambda A. \lambda P \in 2^A. \mathbf{if} (\exists x \in A. p (ap P x)) \mathbf{then} (the (\lambda x. x \in A \wedge P x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 18** We define  $c\_2Eiterate\_2Eneutral$  to be  $\lambda A\_27a : \iota. \lambda V0op \in ((A\_27a^{A\_27a})^{A\_27a}). (ap (c\_2Emin$

**Definition 19** We define  $c\_2Eiterate\_2Emonoidal$  to be  $\lambda A\_27a : \iota. \lambda V0op \in ((A\_27a^{A\_27a})^{A\_27a}). (ap (ap c\_2E$

Let  $ty\_2Enum\_2Enum : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Enum\_2Enum \tag{4}$$

Let  $c\_2Earithmetic\_2E\_2A : \iota$  be given. Assume the following.

$$c\_2Earithmetic\_2E\_2A \in ((ty\_2Enum\_2Enum^{ty\_2Enum\_2Enum})^{ty\_2Enum\_2Enum}) \tag{5}$$

**Definition 20** We define  $c\_2Eiterate\_2Esupport$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}). \lambda V$

**Definition 21** We define  $c\_2Ebool\_2ECOND$  to be  $\lambda A\_27a : \iota. (\lambda V0t \in 2. (\lambda V1t1 \in A\_27a. (\lambda V2t2 \in A\_27a. ($

**Definition 22** We define  $c\_2Eiterate\_2EITSET$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0f \in ((A\_27a^{A\_27a})^{A\_27b}). \lambda V$

**Definition 23** We define  $c\_2Eiterate\_2Eiterate$  to be  $\lambda A\_27a : \iota. \lambda A\_27b : \iota. \lambda V0op \in ((A\_27b^{A\_27b})^{A\_27b}). \lambda V$

**Definition 24** We define  $c\_2Eproduct\_2Enproduct$  to be  $\lambda A\_27a : \iota. (ap (c\_2Eiterate\_2Eiterate A\_27a ty\_2Enum$

Assume the following.

$$True \tag{6}$$

Assume the following.

$$\forall A\_27a. nonempty\ A\_27a \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A\_27a. (p\ V0t)) \Leftrightarrow (p\ V0t))) \tag{7}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2. (((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ & (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ & (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \tag{8}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (9)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.((V0x = V0x) \Leftrightarrow True)) \quad (10)$$

Assume the following.

$$\forall A\_27a.nonempty A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (11)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (12)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (13)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x\_27 \in 2.(\forall V2y \in 2.(\forall V3y\_27 \in 2.(((p V0x) \Leftrightarrow (p V1x\_27)) \wedge ((p V1x\_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y\_27)))))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x\_27) \Rightarrow (p V3y\_27)))))) \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall A\_27a.nonempty A\_27a \Rightarrow \forall A\_27b.nonempty A\_27b \Rightarrow ( \\ & \forall V0op \in ((A\_27a^{A\_27a})^{A\_27a}).((p (ap (c\_2Eiterate\_2Emonoidal \\ & A\_27a) V0op)) \Rightarrow (\forall V1f \in (A\_27a^{A\_27b}).(\forall V2s \in (2^{A\_27b}). \\ & (\forall V3t \in (2^{A\_27b}).(((p (ap (c\_2Epred\_set\_2EFINITE A\_27b) \\ & V2s)) \wedge ((p (ap (c\_2Epred\_set\_2EFINITE A\_27b) V3t)) \wedge (p (ap (ap \\ & (c\_2Epred\_set\_2EDISJOINT A\_27b) V2s) V3t)))))) \Rightarrow ((ap (ap (ap (c\_2Eiterate\_2Eiterate \\ & A\_27b A\_27a) V0op) (ap (ap (c\_2Epred\_set\_2EUNION A\_27b) V2s) V3t)) \\ & V1f) = (ap (ap V0op (ap (ap (ap (c\_2Eiterate\_2Eiterate A\_27b A\_27a) \\ & V0op) V2s) V1f)) (ap (ap (c\_2Eiterate\_2Eiterate A\_27b A\_27a) \\ & V0op) V3t) V1f)))))) \quad (15) \end{aligned}$$

Assume the following.

$$(p (ap (c\_2Eiterate\_2Emonoidal ty\_2Enum\_2Enum) c\_2Earithmetic\_2E\_2A)) \quad (16)$$

**Theorem 1**

$$\begin{aligned} & \forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow (\forall V0f \in (ty\_2Enum\_2Enum^{A_{27a}}). \\ & (\forall V1s \in (2^{A_{27a}}). (\forall V2t \in (2^{A_{27a}}). (((p (ap (c\_2Epred\_set\_2EFINITE \\ & A_{27a}) V1s)) \wedge (p (ap (c\_2Epred\_set\_2EFINITE A_{27a}) V2t)) \wedge (p \\ & (ap (ap (c\_2Epred\_set\_2EDISJOINT A_{27a}) V1s) V2t)))) \Rightarrow ((ap (ap \\ & (c\_2Eproduct\_2Enproduct A_{27a}) (ap (ap (c\_2Epred\_set\_2EUNION \\ & A_{27a}) V1s) V2t)) V0f) = (ap (ap c\_2Earithmetic\_2E\_2A (ap (ap (c\_2Eproduct\_2Enproduct \\ & A_{27a}) V1s) V0f)) (ap (ap (c\_2Eproduct\_2Enproduct A_{27a}) V2t) V0f)))))) \end{aligned}$$