

thm_2EquantHeuristics_2EGUESS_EXISTS_POINT_THM
(TMYhLkA-
NiE8Z6jem1ufweYruiMoZqpVGWGz)

October 26, 2020

Definition 1 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$).

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap V0P (ap (c_2Emin_2E_40 A$

Definition 4 We define $c_2Ebool_2E_T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V$

Definition 5 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$
of type ι .

Definition 6 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A-27a}).(ap (ap (c_2Emin_2E_3D (2^{A-27$

Definition 7 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 8 We define $c_2EquantHeuristics_2EGUESS_EXISTS_POINT$ to be
 $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0i \in (A_27b^{A-27a}).\lambda V1P \in (2^{A-27b}).(ap (c_2Ebool_2E_21 A_27a) (\lambda V2fv \in A_27a.$

Definition 9 We define $c_2Ebool_2E_F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 10 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0f \in (2^{A_{.27a}}).((p\ (ap\ (c_{.2Ebool}._{2E}_{.3F}\ A_{.27a})\ V0f)) \Leftrightarrow (\exists V1x \in A_{.27a}.(p\ (ap\ V0f\ V1x)))))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge (((p\ V0t) \Rightarrow (p\ V0t)) \Leftrightarrow True) \wedge ((p\ V0t) \Rightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (4)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (5)$$

Assume the following.

$$\forall A_{.27a}.nonempty\ A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}.(\forall V1y \in A_{.27a}.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (6)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \quad (7)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2) \Rightarrow (p\ V2t3)) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \quad (8)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x_{.27} \in 2.(\forall V2y \in 2.(\forall V3y_{.27} \in 2.(((p\ V0x) \Leftrightarrow (p\ V1x_{.27})) \wedge ((p\ V1x_{.27}) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y_{.27}))) \Rightarrow (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x_{.27}) \Rightarrow (p\ V3y_{.27})))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \quad (10)$$

Assume the following.

$$(\forall V0A \in 2.((p\ V0A) \Rightarrow ((\neg(p\ V0A)) \Rightarrow False))) \quad (11)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p\ V0A)) \Rightarrow False) \Rightarrow (((p\ V0A) \Rightarrow False) \Rightarrow False))) \quad (12)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p\ V0p) \Rightarrow (p\ V1q))) \Rightarrow (p\ V0p)))) \quad (13)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p\ V0p) \Rightarrow (p\ V1q))) \Rightarrow (\neg(p\ V1q)))) \quad (14)$$

Theorem 1

$$\begin{aligned} & \forall A_{27a}. \text{nonempty } A_{27a} \Rightarrow \forall A_{27b}. \text{nonempty } A_{27b} \Rightarrow (\\ & \quad \forall V0i \in (A_{27b}^{A_{27a}}). (\forall V1P \in (2^{A_{27b}}). ((p \text{ (ap (ap} \\ & \text{(c_2EquantHeuristics_2EGUESS_EXISTS_POINT } A_{27a} \ A_{27b}) \ V0i) \\ & \quad V1P)) \Rightarrow ((p \text{ (ap (c_2Ebool_2E_3F } A_{27b}) \ V1P)) \Leftrightarrow \text{True})))) \end{aligned}$$