

thm_2EquantHeuristics_2EGUESS_POINT_THM (TMPyCLDijgjPvonMYrhwmKTH2QqeMi7bjvy)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 3 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$
of type $\iota \Rightarrow \iota$.

Definition 4 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A))))$

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p P \Rightarrow p Q)$
of type ι .

Definition 6 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 7 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V2t \in 2. V2t)))$

Definition 8 We define `c_2EquantHeuristics_2EGUESS_EXISTS_POINT` to be
 $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a)) (\lambda V2fv \in A. 27a.$

Definition 9 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2)) (\lambda V0t \in 2. V0t)$.

Definition 10 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t)) (\text{c_2Ebool_2E_2F } V0t)))$

Definition 11 We define `c_2EquantHeuristics_2EGUESS_FORALL_POINT` to
be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a)) (\lambda V2fv \in A. 27a.$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (3)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0x \in A.27a.(\forall V1y \in A.27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (6)$$

Assume the following.

$$\forall A.27a.nonempty A.27a \Rightarrow (\forall V0P \in (2^{A.27a}).((\neg(\forall V1x \in A.27a.(p (ap V0P V1x)))) \Leftrightarrow (\exists V2x \in A.27a.(\neg(p (ap V0P V2x)))))) \quad (7)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \quad (8)$$

Assume the following.

$$(\forall V0x \in 2.(\forall V1x.27 \in 2.(\forall V2y \in 2.(\forall V3y.27 \in 2.(((p V0x) \Leftrightarrow (p V1x.27)) \wedge ((p V1x.27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y.27)))) \Rightarrow (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x.27) \Rightarrow (p V3y.27)))))) \quad (9)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (10)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (11)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (12)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (13)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (14)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & \quad \forall V0i \in (A_27b^{A_27a}). (\forall V1P \in (2^{A_27b}). (((p\ (ap\ (ap \\ (c_2EquantHeuristics_2EGUESS_EXISTS_POINT\ A_27a\ A_27b)\ V0i) \\ V1P)) \Rightarrow ((\exists V2v \in A_27b. (p\ (ap\ V1P\ V2v))) \Leftrightarrow True)) \wedge ((p\ (ap\ (ap \\ (c_2EquantHeuristics_2EGUESS_FORALL_POINT\ A_27a\ A_27b)\ V0i) \\ V1P)) \Rightarrow ((\forall V3v \in A_27b. (p\ (ap\ V1P\ V3v))) \Leftrightarrow False)))))) \end{aligned}$$