

# thm\_2EquantHeuristics\_2EGUESS\_RULES\_EQUATION\_EXISTS (TMG4EPDJS7ynwtYhHDCh7wxd8PSmp7AwFH8)

October 26, 2020

Let  $ty\_2Eone\_2Eone : \iota$  be given. Assume the following.

$$nonempty\ ty\_2Eone\_2Eone \quad (1)$$

**Definition 1** We define  $c\_2Emin\_2E3D$  to be  $\lambda A.\lambda x \in A.\lambda y \in A.inj\_o (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define  $c\_2Ebool\_2E2T$  to be  $(ap (ap (c\_2Emin\_2E3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define  $c\_2Ebool\_2E21$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap (ap (c\_2Emin\_2E3D (2^{A\_27a}))$

**Definition 4** We define  $c\_2Ebool\_2E2F$  to be  $(ap (c\_2Ebool\_2E21 2) (\lambda V0t \in 2.V0t))$ .

**Definition 5** We define  $c\_2Emin\_2E3D\_3D\_3E$  to be  $\lambda P \in 2.\lambda Q \in 2.inj\_o (p \Rightarrow q)$  of type  $\iota$ .

**Definition 6** We define  $c\_2Ebool\_2E5C\_2F$  to be  $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c\_2Ebool\_2E21 2) (\lambda V2t \in 2.V2t))$

**Definition 7** We define  $c\_2Emin\_2E40$  to be  $\lambda A.\lambda P \in 2^A.if (\exists x \in A.p (ap P x))$  then (the  $(\lambda x.x \in A \wedge p$  of type  $\iota \Rightarrow \iota$ .

**Definition 8** We define  $c\_2Ebool\_2E3F$  to be  $\lambda A\_27a : \iota.(\lambda V0P \in (2^{A\_27a}).(ap V0P (ap (c\_2Emin\_2E40 A$

**Definition 9** We define  $c\_2Ebool\_2E7E$  to be  $(\lambda V0t \in 2.(ap (ap c\_2Emin\_2E3D\_3D\_3E V0t) c\_2Ebool\_2E2F$

**Definition 10** We define  $c\_2EquantHeuristics\_2EGUESS\_FORALL\_GAP$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0i \in (A\_27b^{A\_27a}).\lambda V1P \in (2^{A\_27b}).(ap (c\_2Ebool\_2E21 A\_27b) (\lambda V2v \in A\_27b.(ap$

**Definition 11** We define  $c\_2EquantHeuristics\_2EGUESS\_EXISTS\_GAP$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0i \in (A\_27b^{A\_27a}).\lambda V1P \in (2^{A\_27b}).(ap (c\_2Ebool\_2E21 A\_27b) (\lambda V2v \in A\_27b.(ap$

**Definition 12** We define  $c\_2EquantHeuristics\_2EGUESS\_FORALL\_POINT$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0i \in (A\_27b^{A\_27a}).\lambda V1P \in (2^{A\_27b}).(ap (c\_2Ebool\_2E21 A\_27a) (\lambda V2fv \in A\_27b.(ap$

**Definition 13** We define `c_2EquantHeuristics_2EGUESS_EXISTS_POINT` to be  $\lambda A_{.27a} : \iota. \lambda A_{.27b} : \iota. \lambda V0i \in (A_{.27b}^{A_{.27a}}). \lambda V1P \in (2^{A_{.27b}}). (ap (c_2Ebool_2E_21 A_{.27a}) (\lambda V2fv \in A_{.27a}.$

**Definition 14** We define `c_2EquantHeuristics_2EGUESS_FORALL` to be  $\lambda A_{.27a} : \iota. \lambda A_{.27b} : \iota. \lambda V0i \in (A_{.27b}^{A_{.27a}}).$

**Definition 15** We define `c_2EquantHeuristics_2EGUESS_EXISTS` to be  $\lambda A_{.27a} : \iota. \lambda A_{.27b} : \iota. \lambda V0i \in (A_{.27b}^{A_{.27a}}).$

**Definition 16** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Assume the following.

$$True \tag{2}$$

Assume the following.

$$\forall A_{.27a}. nonempty A_{.27a} \Rightarrow (\forall V0t \in 2. ((\forall V1x \in A_{.27a}. (p V0t)) \Leftrightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A_{.27a}. nonempty A_{.27a} \Rightarrow (\forall V0t \in 2. ((\exists V1x \in A_{.27a}. (p V0t)) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$\forall A_{.27a}. nonempty A_{.27a} \Rightarrow (\forall V0x \in A_{.27a}. ((V0x = V0x) \Leftrightarrow True)) \tag{5}$$

Assume the following.

$$(\forall V0t \in 2. (((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow \neg(p V0t)) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow \neg(p V0t)))))) \tag{6}$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((p V0A) \Rightarrow (p V1B)) \Leftrightarrow ((\neg(p V0A)) \vee (p V1B)))))) \tag{7}$$

Assume the following.

$$\forall A_{.27a}. nonempty A_{.27a} \Rightarrow (\forall V0f \in (2^{A_{.27a}}). (\forall V1v \in A_{.27a}. ((\forall V2x \in A_{.27a}. ((V2x = V1v) \Rightarrow (p (ap V0f V2x)))) \Leftrightarrow (p (ap V0f V1v)))))) \tag{8}$$

Assume the following.

$$\begin{aligned}
& \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\
& \quad \forall V0i \in (A\_27b^{A\_27a}). (\forall V1P \in (2^{A\_27b}). (((p\ (ap\ ( \\
& \quad ap\ (c\_2EquantHeuristics\_2EGUESS\_EXISTS\ A\_27a\ A\_27b)\ V0i)\ V1P)) \Leftrightarrow \\
& \quad (\forall V2v \in A\_27b. ((p\ (ap\ V1P\ V2v)) \Rightarrow (\exists V3fv \in A\_27a. (p\ ( \\
& \quad ap\ V1P\ (ap\ V0i\ V3fv)))))) \wedge ((p\ (ap\ (ap\ (c\_2EquantHeuristics\_2EGUESS\_FORALL \\
& \quad A\_27a\ A\_27b)\ V0i)\ V1P)) \Leftrightarrow (\forall V4v \in A\_27b. ((\neg(p\ (ap\ V1P\ V4v))) \Rightarrow \\
& \quad (\exists V5fv \in A\_27a. (\neg(p\ (ap\ V1P\ (ap\ V0i\ V5fv)))))) \wedge ((\forall V6i \in \\
& \quad (A\_27b^{A\_27a}). (\forall V7P \in (2^{A\_27b}). ((p\ (ap\ (ap\ (c\_2EquantHeuristics\_2EGUESS\_EXISTS\_POINT \\
& \quad A\_27a\ A\_27b)\ V6i)\ V7P)) \Leftrightarrow (\forall V8fv \in A\_27a. (p\ (ap\ V7P\ (ap\ V6i\ V8fv)))))) \wedge \\
& \quad ((\forall V9i \in (A\_27b^{A\_27a}). (\forall V10P \in (2^{A\_27b}). ((p\ (ap \\
& \quad (ap\ (c\_2EquantHeuristics\_2EGUESS\_FORALL\_POINT\ A\_27a\ A\_27b) \\
& \quad V9i)\ V10P)) \Leftrightarrow (\forall V11fv \in A\_27a. (\neg(p\ (ap\ V10P\ (ap\ V9i\ V11fv)))))) \wedge \\
& \quad ((\forall V12i \in (A\_27b^{A\_27a}). (\forall V13P \in (2^{A\_27b}). ((p\ (ap \\
& \quad (ap\ (c\_2EquantHeuristics\_2EGUESS\_EXISTS\_GAP\ A\_27a\ A\_27b) \\
& \quad V12i)\ V13P)) \Leftrightarrow (\forall V14v \in A\_27b. ((p\ (ap\ V13P\ V14v)) \Rightarrow (\exists V15fv \in \\
& \quad A\_27a. (V14v = (ap\ V12i\ V15fv)))))) \wedge (\forall V16i \in (A\_27b^{A\_27a}). \\
& \quad (\forall V17P \in (2^{A\_27b}). ((p\ (ap\ (ap\ (c\_2EquantHeuristics\_2EGUESS\_FORALL\_GAP \\
& \quad A\_27a\ A\_27b)\ V16i)\ V17P)) \Leftrightarrow (\forall V18v \in A\_27b. ((\neg(p\ (ap\ V17P\ V18v)) \Rightarrow \\
& \quad (\exists V19fv \in A\_27a. (V18v = (ap\ V16i\ V19fv))))))))))))) \\
& \hspace{15em} (9)
\end{aligned}$$

**Theorem 1**

$$\begin{aligned}
& \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0i \in A\_27a. (p\ (ap\ (ap\ (c\_2EquantHeuristics\_2EGUESS\_EXISTS\_GAP \\
& \quad ty\_2Eone\_2Eone\ A\_27a)\ (\lambda V1xxx \in ty\_2Eone\_2Eone.V0i))\ (\lambda V2x \in \\
& \quad A\_27a. (ap\ (ap\ (c\_2Emin\_2E\_3D\ A\_27a)\ V2x)\ V0i))))))
\end{aligned}$$