

thm_2EquantHeuristics_2EGUESS__RULES__EXISTS__UNIQUE (TMYhH2C1ewsyoEgePHW7tndmfDdL6oFUpuP)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 3 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))$

Definition 4 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2. V0t))$.

Definition 5 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$
of type ι .

Definition 6 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))$

Definition 7 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p (\text{ap } P x)))$
of type $\iota \Rightarrow \iota$.

Definition 8 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A$

Definition 9 We define `c_2Ebool_2E_3F_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } \text{c_2Ebool_2E_2F_5C } (2^{A-27a}))$

Definition 10 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } \text{c_2Emin_2E_3D_3D_3E } V0t) \text{ c_2Ebool_2E_2F_5C } (2^{A-27a}))$

Definition 11 We define `c_2EquantHeuristics_2EGUESS__FORALL__GAP` to be
 $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b) (\lambda V2v \in A. 27b. ($

Definition 12 We define `c_2EquantHeuristics_2EGUESS__EXISTS__GAP` to be
 $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b) (\lambda V2v \in A. 27b. ($

Definition 13 We define `c_2EquantHeuristics_2EGUESS__FORALL__POINT` to
be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a) (\lambda V2fv \in A. 27a. ($

Definition 14 We define `c_2EquantHeuristics_2EGUESS__EXISTS__POINT` to be
 $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a) (\lambda V2fv \in A. 27a. ($

Definition 15 We define `c_2EquantHeuristics_2EGUESS_FORALL` to be $\lambda A_{.27a} : \iota.\lambda A_{.27b} : \iota.\lambda V0i \in (A_{.27a} \dots$

Definition 16 We define `c_2EquantHeuristics_2EGUESS_EXISTS` to be $\lambda A_{.27a} : \iota.\lambda A_{.27b} : \iota.\lambda V0i \in (A_{.27a} \dots$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p V0t))) \tag{3}$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0t \in 2.((\forall V1x \in A_{.27a}.(p V0t)) \Leftrightarrow (p V0t))) \tag{4}$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0t \in 2.((\exists V1x \in A_{.27a}.(p V0t)) \Leftrightarrow (p V0t))) \tag{5}$$

Assume the following.

$$(\forall V0t \in 2.(((True \wedge (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \wedge True) \Leftrightarrow (p V0t)) \wedge (((False \wedge (p V0t)) \Leftrightarrow False) \wedge (((p V0t) \wedge False) \Leftrightarrow False) \wedge (((p V0t) \wedge (p V0t)) \Leftrightarrow (p V0t)))))) \tag{6}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{7}$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)))) \tag{8}$$

Assume the following.

$$\forall A_{.27a}.nonempty A_{.27a} \Rightarrow (\forall V0x \in A_{.27a} . (\forall V1y \in A_{.27a} . ((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \tag{9}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \tag{10}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). (\forall V1Q \in \\ & 2. ((\forall V2x \in A_27a. ((p\ (ap\ V0P\ V2x)) \Rightarrow (p\ V1Q))) \Leftrightarrow ((\exists V3x \in \\ & A_27a. (p\ (ap\ V0P\ V3x)) \Rightarrow (p\ V1Q)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2. (\forall V1t2 \in 2. (\forall V2t3 \in 2. (((p\ V0t1) \Rightarrow \\ & ((p\ V1t2) \Rightarrow (p\ V2t3))) \Leftrightarrow (((p\ V0t1) \wedge (p\ V1t2)) \Rightarrow (p\ V2t3)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow (\forall V0P \in (2^{A_27a}). ((p\ (ap \\ & (c_2Ebool_2E_3F_21\ A_27a)\ (\lambda V1x \in A_27a. (ap\ V0P\ V1x)))) \Leftrightarrow ((\\ & \exists V2x \in A_27a. (p\ (ap\ V0P\ V2x))) \wedge (\forall V3x \in A_27a. (\forall V4y \in \\ & A_27a. (((p\ (ap\ V0P\ V3x)) \wedge (p\ (ap\ V0P\ V4y))) \Rightarrow (V3x = V4y)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2. (\forall V1x_27 \in 2. (\forall V2y \in 2. (\forall V3y_27 \in \\ & 2. (((((p\ V0x) \Leftrightarrow (p\ V1x_27)) \wedge ((p\ V1x_27) \Rightarrow ((p\ V2y) \Leftrightarrow (p\ V3y_27)))) \Rightarrow \\ & (((p\ V0x) \Rightarrow (p\ V2y)) \Leftrightarrow ((p\ V1x_27) \Rightarrow (p\ V3y_27)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\ & \forall V0i \in (A_27b^{A_27a}). (\forall V1P \in (2^{A_27b}). (((p\ (ap\ (\\ & ap\ (c_2EquantHeuristics_2EGUESS_EXISTS\ A_27a\ A_27b)\ V0i)\ V1P)) \Leftrightarrow \\ & (\forall V2v \in A_27b. ((p\ (ap\ V1P\ V2v)) \Rightarrow (\exists V3fv \in A_27a. (p\ (\\ & ap\ V1P\ (ap\ V0i\ V3fv)))))) \wedge ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL \\ & A_27a\ A_27b)\ V0i)\ V1P)) \Leftrightarrow (\forall V4v \in A_27b. ((\neg (p\ (ap\ V1P\ V4v))) \Rightarrow \\ & (\exists V5fv \in A_27a. (\neg (p\ (ap\ V1P\ (ap\ V0i\ V5fv)))))) \wedge ((\forall V6i \in \\ & (A_27b^{A_27a}). (\forall V7P \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_POINT \\ & A_27a\ A_27b)\ V6i)\ V7P)) \Leftrightarrow (\forall V8fv \in A_27a. (p\ (ap\ V7P\ (ap\ V6i\ V8fv)))))) \wedge \\ & ((\forall V9i \in (A_27b^{A_27a}). (\forall V10P \in (2^{A_27b}). ((p\ (ap \\ & (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_POINT\ A_27a\ A_27b)\ \\ & V9i)\ V10P)) \Leftrightarrow (\forall V11fv \in A_27a. (\neg (p\ (ap\ V10P\ (ap\ V9i\ V11fv)))))) \wedge \\ & ((\forall V12i \in (A_27b^{A_27a}). (\forall V13P \in (2^{A_27b}). ((p\ (ap \\ & (ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_GAP\ A_27a\ A_27b)\ \\ & V12i)\ V13P)) \Leftrightarrow (\forall V14v \in A_27b. ((p\ (ap\ V13P\ V14v)) \Rightarrow (\exists V15fv \in \\ & A_27a. (V14v = (ap\ V12i\ V15fv)))))) \wedge (\forall V16i \in (A_27b^{A_27a}). \\ & (\forall V17P \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_GAP \\ & A_27a\ A_27b)\ V16i)\ V17P)) \Leftrightarrow (\forall V18v \in A_27b. ((\neg (p\ (ap\ V17P\ V18v))) \Rightarrow \\ & (\exists V19fv \in A_27a. (V18v = (ap\ V16i\ V19fv)))))) \end{aligned} \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow \forall A_27c. \\ & nonempty\ A_27c \Rightarrow (\forall V0i \in (A_27c^{A_27b}). (\forall V1P \in ((2^{A_27a})^{A_27c}). \\ & (((\forall V2y \in A_27a.(p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_POINT \\ & A_27b\ A_27c)\ V0i)\ (\lambda V3x \in A_27c.(ap\ (ap\ V1P\ V3x)\ V2y)))))) \Rightarrow (p\ (\\ & ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_POINT\ A_27b\ A_27c) \\ & V0i)\ (\lambda V4x \in A_27c.(ap\ (c_2Ebool_2E_3F_21\ A_27a)\ (\lambda V5y \in \\ & A_27a.(ap\ (ap\ V1P\ V4x)\ V5y)))))) \wedge ((\forall V6y \in A_27a.(p\ (ap\ (\\ & ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_GAP\ A_27b\ A_27c)\ V0i) \\ & (\lambda V7x \in A_27c.(ap\ (ap\ V1P\ V7x)\ V6y)))))) \Rightarrow (p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_GAP \\ & A_27b\ A_27c)\ V0i)\ (\lambda V8x \in A_27c.(ap\ (c_2Ebool_2E_3F_21\ A_27a) \\ & (\lambda V9y \in A_27a.(ap\ (ap\ V1P\ V8x)\ V9y)))))))))) \end{aligned}$$