

thm_2EquantHeuristics_2EGUESS_RULES_WEAKEN_FORALL (TMKJvty3TKWdh7NDSwEXK16YdhB1HJ8dPAK)

October 26, 2020

Definition 1 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 2 We define `c_2Ebool_2ET` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 3 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (\text{the } (\lambda x. x \in A \wedge p x))$
of type $\iota \Rightarrow \iota$.

Definition 4 We define `c_2Ebool_2E_3F` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c_2Emin_2E_40 } A))))$

Definition 5 We define `c_2Ebool_2E_21` to be $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A-27a}))))$

Definition 6 We define `c_2Ebool_2EF` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2.V0t))$.

Definition 7 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$
of type ι .

Definition 8 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) (\text{c_2Ebool_2EF } 2))))$

Definition 9 We define `c_2EquantHeuristics_2EGUESS_FORALL_GAP` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b)) (\lambda V2v \in A. 27b. (p \Rightarrow q))$

Definition 10 We define `c_2EquantHeuristics_2EGUESS_EXISTS_GAP` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b)) (\lambda V2v \in A. 27b. (p \Rightarrow q))$

Definition 11 We define `c_2EquantHeuristics_2EGUESS_FORALL_POINT` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a)) (\lambda V2fv \in A. 27a. (p \Rightarrow q))$

Definition 12 We define `c_2EquantHeuristics_2EGUESS_EXISTS_POINT` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). \lambda V1P \in (2^{A-27b}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27a)) (\lambda V2fv \in A. 27a. (p \Rightarrow q))$

Definition 13 We define `c_2EquantHeuristics_2EGUESS_FORALL` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b)) (\lambda V2v \in A. 27b. (p \Rightarrow q))$

Definition 14 We define `c_2EquantHeuristics_2EGUESS_EXISTS` to be $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0i \in (A. 27b^{A-27a}). (\text{ap } (\text{c_2Ebool_2E_21 } A. 27b)) (\lambda V2v \in A. 27b. (p \Rightarrow q))$

Definition 15 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Definition 16 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow \\ & True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((\\ & (p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & ((\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \wedge ((\neg True) \Leftrightarrow False) \wedge \\ & ((\neg False) \Leftrightarrow True))) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall A_27a.nonempty A_27a \Rightarrow (\forall V0x \in A_27a.(\forall V1y \in \\ & A_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \end{aligned} \tag{4}$$

Assume the following.

$$\begin{aligned} & (\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow \\ & (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(\\ & p V0t)))))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} & (\forall V0t1 \in 2.(\forall V1t2 \in 2.(\forall V2t3 \in 2.(((p V0t1) \Rightarrow \\ & ((p V1t2) \Rightarrow (p V2t3))) \Leftrightarrow (((p V0t1) \wedge (p V1t2)) \Rightarrow (p V2t3)))))) \end{aligned} \tag{6}$$

Assume the following.

$$\begin{aligned} & (\forall V0x \in 2.(\forall V1x_27 \in 2.(\forall V2y \in 2.(\forall V3y_27 \in \\ & 2.(((p V0x) \Leftrightarrow (p V1x_27)) \wedge ((p V1x_27) \Rightarrow ((p V2y) \Leftrightarrow (p V3y_27)))) \Rightarrow \\ & (((p V0x) \Rightarrow (p V2y)) \Leftrightarrow ((p V1x_27) \Rightarrow (p V3y_27)))))) \end{aligned} \tag{7}$$

Assume the following.

$$\begin{aligned}
& \forall A_27a.nonempty\ A_27a \Rightarrow \forall A_27b.nonempty\ A_27b \Rightarrow (\\
& \quad \forall V0i \in (A_27b^{A_27a}). (\forall V1P \in (2^{A_27b}). (((p\ (ap\ (\\
& \quad ap\ (c_2EquantHeuristics_2EGUESS_EXISTS\ A_27a\ A_27b)\ V0i)\ V1P))) \Leftrightarrow \\
& \quad (\forall V2v \in A_27b. ((p\ (ap\ V1P\ V2v)) \Rightarrow (\exists V3fv \in A_27a. (p\ (\\
& \quad ap\ V1P\ (ap\ V0i\ V3fv)))))) \wedge ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL \\
& \quad A_27a\ A_27b)\ V0i)\ V1P))) \Leftrightarrow (\forall V4v \in A_27b. ((\neg(p\ (ap\ V1P\ V4v))) \Rightarrow \\
& \quad (\exists V5fv \in A_27a. (\neg(p\ (ap\ V1P\ (ap\ V0i\ V5fv)))))) \wedge ((\forall V6i \in \\
& \quad (A_27b^{A_27a}). (\forall V7P \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_POINT \\
& \quad A_27a\ A_27b)\ V6i)\ V7P))) \Leftrightarrow (\forall V8fv \in A_27a. (p\ (ap\ V7P\ (ap\ V6i\ V8fv)))))) \wedge \\
& \quad ((\forall V9i \in (A_27b^{A_27a}). (\forall V10P \in (2^{A_27b}). ((p\ (ap \\
& \quad (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_POINT\ A_27a\ A_27b) \\
& \quad V9i)\ V10P))) \Leftrightarrow (\forall V11fv \in A_27a. (\neg(p\ (ap\ V10P\ (ap\ V9i\ V11fv)))))) \wedge \\
& \quad ((\forall V12i \in (A_27b^{A_27a}). (\forall V13P \in (2^{A_27b}). ((p\ (ap \\
& \quad (ap\ (c_2EquantHeuristics_2EGUESS_EXISTS_GAP\ A_27a\ A_27b) \\
& \quad V12i)\ V13P))) \Leftrightarrow (\forall V14v \in A_27b. ((p\ (ap\ V13P\ V14v)) \Rightarrow (\exists V15fv \in \\
& \quad A_27a. (V14v = (ap\ V12i\ V15fv)))))) \wedge (\forall V16i \in (A_27b^{A_27a}). \\
& \quad (\forall V17P \in (2^{A_27b}). ((p\ (ap\ (ap\ (c_2EquantHeuristics_2EGUESS_FORALL_GAP \\
& \quad A_27a\ A_27b)\ V16i)\ V17P))) \Leftrightarrow (\forall V18v \in A_27b. ((\neg(p\ (ap\ V17P\ V18v))) \Rightarrow \\
& \quad (\exists V19fv \in A_27a. (V18v = (ap\ V16i\ V19fv))))))))) \\
& \hspace{10em} (8)
\end{aligned}$$

Assume the following.

$$(\forall V0t \in 2. ((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \quad (9)$$

Assume the following.

$$(\forall V0A \in 2. ((p\ V0A) \Rightarrow ((\neg(p\ V0A)) \Rightarrow False))) \quad (10)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((p\ V0A) \vee (p\ V1B))) \Rightarrow False) \Leftrightarrow ((p\ V0A) \Rightarrow False) \Rightarrow ((\neg(p\ V1B)) \Rightarrow False)))) \quad (11)$$

Assume the following.

$$(\forall V0A \in 2. (\forall V1B \in 2. (((\neg((\neg(p\ V0A)) \vee (p\ V1B))) \Rightarrow False) \Leftrightarrow ((p\ V0A) \Rightarrow ((\neg(p\ V1B)) \Rightarrow False)))) \quad (12)$$

Assume the following.

$$(\forall V0A \in 2. (((\neg(p\ V0A)) \Rightarrow False) \Rightarrow (((p\ V0A) \Rightarrow False) \Rightarrow False))) \quad (13)$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p\ V0p) \Leftrightarrow (\\
& \quad (p\ V1q) \Leftrightarrow (p\ V2r))) \Leftrightarrow (((p\ V0p) \vee ((p\ V1q) \vee (p\ V2r))) \wedge (((p\ V0p) \vee ((\neg \\
& \quad p\ V2r)) \vee (\neg(p\ V1q)))) \wedge (((p\ V1q) \vee ((\neg(p\ V2r)) \vee (\neg(p\ V0p)))) \wedge ((p\ V2r) \vee \\
& \quad ((\neg(p\ V1q)) \vee (\neg(p\ V0p))))))))) \\
& \hspace{10em} (14)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow (\\
& (p \ V1q) \vee (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee \neg(p \ V1q)) \wedge (((p \ V0p) \vee \neg(p \ V2r))) \wedge \\
& ((p \ V1q) \vee ((p \ V2r) \vee \neg(p \ V0p))))))))))
\end{aligned} \tag{15}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (\forall V2r \in 2. (((p \ V0p) \Leftrightarrow (\\
& (p \ V1q) \Rightarrow (p \ V2r))) \Leftrightarrow (((p \ V0p) \vee (p \ V1q)) \wedge (((p \ V0p) \vee \neg(p \ V2r))) \wedge (\\
& \neg(p \ V1q) \vee ((p \ V2r) \vee \neg(p \ V0p))))))))))
\end{aligned} \tag{16}$$

Assume the following.

$$\begin{aligned}
& (\forall V0p \in 2. (\forall V1q \in 2. (((p \ V0p) \Leftrightarrow \neg(p \ V1q)) \Leftrightarrow (((p \ V0p) \vee \\
& (p \ V1q)) \wedge (\neg(p \ V1q) \vee \neg(p \ V0p))))))
\end{aligned} \tag{17}$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow (p \ V0p))) \tag{18}$$

Assume the following.

$$(\forall V0p \in 2. (\forall V1q \in 2. (\neg((p \ V0p) \Rightarrow (p \ V1q))) \Rightarrow \neg(p \ V1q))) \tag{19}$$

Theorem 1

$$\begin{aligned}
& \forall A.27a.nonempty \ A.27a \Rightarrow \forall A.27b.nonempty \ A.27b \Rightarrow (\\
& \forall V0i \in (A.27a^{A.27b}). (\forall V1P \in (2^{A.27a}). (\forall V2Q \in \\
& (2^{A.27a}). ((\forall V3x \in A.27a. ((p \ (ap \ V2Q \ V3x)) \Rightarrow (p \ (ap \ V1P \ V3x)))))) \Rightarrow \\
& ((p \ (ap \ (ap \ (c.2EquantHeuristics_2EGUESS_FORALL_POINT \ A.27b \\
& A.27a) \ V0i) \ V1P)) \Rightarrow (p \ (ap \ (ap \ (c.2EquantHeuristics_2EGUESS_FORALL_POINT \\
& A.27b \ A.27a) \ V0i) \ V2Q))))))
\end{aligned}$$