

# thm\_2EquantHeuristics\_2EINL\_\_NEQ\_\_ELIM (TMVVnA8ehStyD1tMJCXPYUD4YGqAXCUgatg)

October 26, 2020

**Definition 1** We define `c_2Emin_2E_3D` to be  $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj\_o } (x = y)$  of type  $\iota \Rightarrow \iota$ .

**Definition 2** We define `c_2Ebool_2E_2T` to be  $(\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

**Definition 3** We define `c_2Ebool_2E_21` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } (\text{ap } (\text{c\_2Emin\_2E\_3D } (2^{A-27a}))$

**Definition 4** We define `c_2Ebool_2E_2F` to be  $(\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V0t \in 2.V0t)$ .

**Definition 5** We define `c_2Emin_2E_3D_3D_3E` to be  $\lambda P \in 2. \lambda Q \in 2. \text{inj\_o } (p P \Rightarrow p Q)$  of type  $\iota$ .

**Definition 6** We define `c_2Emin_2E_40` to be  $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p (\text{ap } P x)) \text{ then } (the (\lambda x. x \in A \wedge p x))$  of type  $\iota \Rightarrow \iota$ .

**Definition 7** We define `c_2Ebool_2E_3F` to be  $\lambda A. 27a : \iota. (\lambda V0P \in (2^{A-27a}). (\text{ap } V0P (\text{ap } (\text{c\_2Emin\_2E\_40 } A$

**Definition 8** We define `c_2Ebool_2E_5C_2F` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V2t \in 2.V2t))$

Let `ty_2Esum_2Esum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A0. \text{nonempty } A0 \Rightarrow \forall A1. \text{nonempty } A1 \Rightarrow \text{nonempty } (ty\_2Esum\_2Esum A0 A1) \tag{1}$$

**Definition 9** We define `c_2Ebool_2E_2F_5C` to be  $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c\_2Ebool\_2E\_21 } 2)) (\lambda V2t \in 2.V2t))$

Let `c_2Esum_2EABS_sum` :  $\iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall A. 27a. \text{nonempty } A. 27a \Rightarrow \forall A. 27b. \text{nonempty } A. 27b \Rightarrow c\_2Esum\_2EABS\_sum A. 27a A. 27b \in ((ty\_2Esum\_2Esum A. 27a A. 27b)^{((2^{A-27b})^{A-27a})^2}) \tag{2}$$

**Definition 10** We define `c_2Esum_2EINL` to be  $\lambda A. 27a : \iota. \lambda A. 27b : \iota. \lambda V0e \in A. 27a. (\text{ap } (c\_2Esum\_2EABS\_sum$

**Definition 11** We define `c_2Ebool_2E_7E` to be  $(\lambda V0t \in 2. (\text{ap } (\text{ap } c\_2Emin\_2E_3D_3D_3E V0t) c\_2Ebool_2E_2F_5C$

**Definition 12** We define  $c\_2Esum\_2EINR$  to be  $\lambda A\_27a : \iota.\lambda A\_27b : \iota.\lambda V0e \in A\_27b.(ap (c\_2Esum\_2EABS$

Let  $c\_2Esum\_2EISR : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow c\_2Esum\_2EISR \\ A\_27a\ A\_27b \in (2^{(ty\_2Esum\_2Esum\ A\_27a\ A\_27b)}) \end{aligned} \quad (3)$$

Assume the following.

$$True \quad (4)$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p\ V0t1) \Rightarrow (p\ V1t2)) \Rightarrow (((p\ V1t2) \Rightarrow (p\ V0t1)) \Rightarrow ((p\ V0t1) \Leftrightarrow (p\ V1t2)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.(False \Rightarrow (p\ V0t))) \quad (6)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0t \in 2.((\forall V1x \in A\_27a.(p\ V0t) \Leftrightarrow (p\ V0t)))) \quad (7)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \wedge True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \wedge (p\ V0t)) \Leftrightarrow False) \wedge (((p\ V0t) \wedge False) \Leftrightarrow False) \wedge \\ (((p\ V0t) \wedge (p\ V0t)) \Leftrightarrow (p\ V0t)))))) \end{aligned} \quad (8)$$

Assume the following.

$$((\forall V0t \in 2.((\neg(\neg(p\ V0t))) \Leftrightarrow (p\ V0t))) \wedge (((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True))) \quad (9)$$

Assume the following.

$$\forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0x \in A\_27a.(\forall V1y \in A\_27a.((V0x = V1y) \Leftrightarrow (V1y = V0x)))) \quad (10)$$

Assume the following.

$$\begin{aligned} (\forall V0t \in 2.(((True \Leftrightarrow (p\ V0t)) \Leftrightarrow (p\ V0t)) \wedge (((p\ V0t) \Leftrightarrow True) \Leftrightarrow \\ (p\ V0t)) \wedge (((False \Leftrightarrow (p\ V0t)) \Leftrightarrow (\neg(p\ V0t))) \wedge (((p\ V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p\ V0t)))))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow (\forall V0f \in (2^{A\_27a}).(\forall V1v \in A\_27a.((\forall V2x \in A\_27a.((V2x = V1v) \Rightarrow (p\ (ap\ V0f\ V2x))) \Leftrightarrow (p\ (ap\ V0f\ V1v)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ (\forall V0y \in A\_27a. (\forall V1x \in A\_27a. (((ap\ (c\_2Esum\_2EINL \\ A\_27a\ A\_27b)\ V1x) = (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V0y))) \Leftrightarrow (V1x = \\ V0y)))) \wedge (\forall V2y \in A\_27b. (\forall V3x \in A\_27b. (((ap\ (c\_2Esum\_2EINR \\ A\_27a\ A\_27b)\ V3x) = (ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ V2y))) \Leftrightarrow (V3x = \\ V2y)))))) \end{aligned} \quad (13)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ \forall V0ss \in (ty\_2Esum\_2Esum\ A\_27a\ A\_27b). ((\exists V1x \in A\_27a. \\ (V0ss = (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V1x))) \vee (\exists V2y \in A\_27b. \\ (V0ss = (ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ V2y)))))) \end{aligned} \quad (14)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ \forall V0x \in A\_27a. (\forall V1y \in A\_27b. (\neg((ap\ (c\_2Esum\_2EINL \\ A\_27a\ A\_27b)\ V0x) = (ap\ (c\_2Esum\_2EINR\ A\_27a\ A\_27b)\ V1y)))))) \end{aligned} \quad (15)$$

Assume the following.

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ (\forall V0x \in A\_27b. (p\ (ap\ (c\_2Esum\_2EISR\ A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EINR \\ A\_27a\ A\_27b)\ V0x)))) \wedge (\forall V1y \in A\_27a. (\neg(p\ (ap\ (c\_2Esum\_2EISR \\ A\_27a\ A\_27b)\ (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V1y)))))) \end{aligned} \quad (16)$$

**Theorem 1**

$$\begin{aligned} \forall A\_27a.nonempty\ A\_27a \Rightarrow \forall A\_27b.nonempty\ A\_27b \Rightarrow ( \\ \forall V0x \in (ty\_2Esum\_2Esum\ A\_27a\ A\_27b). (((\forall V1l \in A\_27a. \\ (\neg(V0x = (ap\ (c\_2Esum\_2EINL\ A\_27a\ A\_27b)\ V1l)))) \Leftrightarrow (p\ (ap\ (c\_2Esum\_2EISR \\ A\_27a\ A\_27b)\ V0x)))) \wedge ((\forall V2l \in A\_27a. (\neg((ap\ (c\_2Esum\_2EINL \\ A\_27a\ A\_27b)\ V2l) = V0x)))) \Leftrightarrow (p\ (ap\ (c\_2Esum\_2EISR\ A\_27a\ A\_27b)\ V0x)))) \end{aligned}$$