

thm_2EquantHeuristics_2EIS_REMOVABLE_QUANT_FUN_____

(TMYSXRpTFvJzDXHaEtS-
gZbZ3W48puFJGaVN)

October 26, 2020

Definition 1 We define $c_2Emin_2E_40$ to be $\lambda A.\lambda P \in 2^A.$ if $(\exists x \in A.p (ap P x))$ then (the $(\lambda x.x \in A \wedge p$
of type $\iota \Rightarrow \iota$).

Definition 2 We define $c_2Emin_2E_3D$ to be $\lambda A.\lambda x \in A.\lambda y \in A.inj_o (x = y)$
of type $\iota \Rightarrow \iota$.

Definition 3 We define $c_2Ebool_2E_2T$ to be $(ap (ap (c_2Emin_2E_3D (2^2)) (\lambda V0x \in 2.V0x)) (\lambda V1x \in 2.V1x))$

Definition 4 We define $c_2Ebool_2E_3F$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap V0P (ap (c_2Emin_2E_40 A_27a$

Definition 5 We define $c_2Ebool_2E_21$ to be $\lambda A_27a : \iota.(\lambda V0P \in (2^{A_27a}).(ap (ap (c_2Emin_2E_3D (2^{A_27a}$

Definition 6 We define $c_2EquantHeuristics_2EIS_REMOVABLE_QUANT_FUN$
to be $\lambda A_27a : \iota.\lambda A_27b : \iota.\lambda V0f \in (A_27b^{A_27a}).(ap (c_2Ebool_2E_21 A_27b) (\lambda V1v \in A_27b.(ap (c_2Ebool$

Definition 7 We define $c_2Ebool_2E_2F$ to be $(ap (c_2Ebool_2E_21 2) (\lambda V0t \in 2.V0t))$.

Definition 8 We define $c_2Emin_2E_3D_3D_3E$ to be $\lambda P \in 2.\lambda Q \in 2.inj_o (p P \Rightarrow p Q)$
of type ι .

Definition 9 We define $c_2Ebool_2E_2F_5C$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 10 We define $c_2Ebool_2E_5C_2F$ to be $(\lambda V0t1 \in 2.(\lambda V1t2 \in 2.(ap (c_2Ebool_2E_21 2) (\lambda V2t \in 2$

Definition 11 We define $c_2Ebool_2E_7E$ to be $(\lambda V0t \in 2.(ap (ap c_2Emin_2E_3D_3D_3E V0t) c_2Ebool_2E$

Assume the following.

$$True \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2.(\forall V1t2 \in 2.(((p V0t1) \Rightarrow (p V1t2)) \Rightarrow (((p V1t2) \Rightarrow (p V0t1)) \Rightarrow ((p V0t1) \Leftrightarrow (p V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))))) \quad (8)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))))) \quad (9)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (10)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee (\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (11)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (12)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (13)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (14)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V0p)))) \quad (15)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V1q)))) \quad (16)$$

Assume the following.

$$(\forall V0p \in 2.((\neg(\neg(p V0p))) \Rightarrow (p V0p))) \quad (17)$$

Theorem 1

$$\begin{aligned} & \forall A.27a.nonempty \ A.27a \Rightarrow \forall A.27b.nonempty \ A.27b \Rightarrow (\\ \forall V0f \in (A.27b^{A.27a}).(\forall V1P \in (2^{A.27b}).((p \ (ap \ (c.2EquantHeuristics.2EIS.REMOVABLE.QUANT.FUN \\ A.27a \ A.27b) \ V0f)) \Rightarrow ((\exists V2x \in A.27a.(p \ (ap \ V1P \ (ap \ V0f \ V2x)))) \Leftrightarrow \\ & (\exists V3x.27 \in A.27b.(p \ (ap \ V1P \ V3x.27)))))) \end{aligned}$$