

thm_2EquantHeuristics_2EIS_REMOVABLE_QUANT_FUN_____

(TMdzDRqov2i6vHSxhkj1Ns4AX1zY7XXdrDa)

October 26, 2020

Definition 1 We define `c_2Emin_2E_40` to be $\lambda A. \lambda P \in 2^A. \text{if } (\exists x \in A. p \text{ (ap } P \ x)) \text{ then (the } (\lambda x. x \in A \wedge p \ x) \text{ of type } \iota \Rightarrow \iota.$

Definition 2 We define `c_2Emin_2E_3D` to be $\lambda A. \lambda x \in A. \lambda y \in A. \text{inj_o } (x = y)$ of type $\iota \Rightarrow \iota.$

Definition 3 We define `c_2Ebool_2E_2T` to be $(\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^2)) (\lambda V0x \in 2. V0x)) (\lambda V1x \in 2. V1x))$

Definition 4 We define `c_2Ebool_2E_3F` to be $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (\text{ap } V0P \ (\text{ap } (\text{c_2Emin_2E_40 } A_{27a}))))$

Definition 5 We define `c_2Ebool_2E_21` to be $\lambda A_{27a} : \iota. (\lambda V0P \in (2^{A_{27a}}). (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D } (2^{A_{27a}}))))$

Definition 6 We define `c_2EquantHeuristics_2EIS_REMOVABLE_QUANT_FUN` to be $\lambda A_{27a} : \iota. \lambda A_{27b} : \iota. \lambda V0f \in (A_{27b}^{A_{27a}}). (\text{ap } (\text{c_2Ebool_2E_21 } A_{27b}) (\lambda V1v \in A_{27b}. (\text{ap } (\text{c_2Ebool_2E_21 } v))))$

Definition 7 We define `c_2Ebool_2E_2F` to be $(\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V0t \in 2. V0t)).$

Definition 8 We define `c_2Emin_2E_3D_3D_3E` to be $\lambda P \in 2. \lambda Q \in 2. \text{inj_o } (p \Rightarrow q)$ of type $\iota.$

Definition 9 We define `c_2Ebool_2E_2F_5C` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))))$

Definition 10 We define `c_2Ebool_2E_5C_2F` to be $(\lambda V0t1 \in 2. (\lambda V1t2 \in 2. (\text{ap } (\text{c_2Ebool_2E_21 } 2) (\lambda V2t \in 2. V2t))))$

Definition 11 We define `c_2Ebool_2E_7E` to be $(\lambda V0t \in 2. (\text{ap } (\text{ap } (\text{c_2Emin_2E_3D_3D_3E } V0t) (\text{c_2Ebool_2E_21 } 2))))$

Assume the following.

$$\text{True} \tag{1}$$

Assume the following.

$$(\forall V0t1 \in 2. (\forall V1t2 \in 2. (((p \ V0t1) \Rightarrow (p \ V1t2)) \Rightarrow (((p \ V1t2) \Rightarrow (p \ V0t1)) \Rightarrow ((p \ V0t1) \Leftrightarrow (p \ V1t2)))))) \tag{2}$$

Assume the following.

$$(\forall V0t \in 2.(((True \Rightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Rightarrow True) \Leftrightarrow True) \wedge (((False \Rightarrow (p V0t)) \Leftrightarrow True) \wedge (((p V0t) \Rightarrow (p V0t)) \Leftrightarrow True) \wedge ((p V0t) \Rightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (3)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t)) \wedge ((\neg True) \Leftrightarrow False) \wedge ((\neg False) \Leftrightarrow True)) \quad (4)$$

Assume the following.

$$(\forall V0t \in 2.(((True \Leftrightarrow (p V0t)) \Leftrightarrow (p V0t)) \wedge (((p V0t) \Leftrightarrow True) \Leftrightarrow (p V0t)) \wedge (((False \Leftrightarrow (p V0t)) \Leftrightarrow (\neg(p V0t))) \wedge (((p V0t) \Leftrightarrow False) \Leftrightarrow (\neg(p V0t)))))) \quad (5)$$

Assume the following.

$$(\forall V0t \in 2.((\neg(\neg(p V0t))) \Leftrightarrow (p V0t))) \quad (6)$$

Assume the following.

$$(\forall V0A \in 2.((p V0A) \Rightarrow ((\neg(p V0A)) \Rightarrow False))) \quad (7)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((p V0A) \vee (p V1B))) \Rightarrow False) \Leftrightarrow (((p V0A) \Rightarrow False) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (8)$$

Assume the following.

$$(\forall V0A \in 2.(\forall V1B \in 2.(((\neg((\neg(p V0A)) \vee (p V1B))) \Rightarrow False) \Leftrightarrow ((p V0A) \Rightarrow ((\neg(p V1B)) \Rightarrow False)))) \quad (9)$$

Assume the following.

$$(\forall V0A \in 2.(((\neg(p V0A)) \Rightarrow False) \Rightarrow (((p V0A) \Rightarrow False) \Rightarrow False))) \quad (10)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Leftrightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee ((p V1q) \vee (p V2r))) \wedge (((p V0p) \vee (\neg(p V2r)) \vee (\neg(p V1q)))) \wedge (((p V1q) \vee ((\neg(p V2r)) \vee (\neg(p V0p)))) \wedge ((p V2r) \vee ((\neg(p V1q)) \vee (\neg(p V0p)))))))))) \quad (11)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.(\forall V2r \in 2.(((p V0p) \Leftrightarrow ((p V1q) \Rightarrow (p V2r))) \Leftrightarrow (((p V0p) \vee (p V1q)) \wedge (((p V0p) \vee (\neg(p V2r))) \wedge ((\neg(p V1q)) \vee ((p V2r) \vee (\neg(p V0p)))))))))) \quad (12)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (p V0p)))) \quad (13)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \Rightarrow (p V1q))) \Rightarrow (\neg(p V1q))))) \quad (14)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V0p))))) \quad (15)$$

Assume the following.

$$(\forall V0p \in 2.(\forall V1q \in 2.((\neg((p V0p) \vee (p V1q))) \Rightarrow (\neg(p V1q))))) \quad (16)$$

Assume the following.

$$(\forall V0p \in 2.((\neg(\neg(p V0p))) \Rightarrow (p V0p))) \quad (17)$$

Theorem 1

$$\begin{aligned} & \forall A.27a.nonempty \ A.27a \Rightarrow \forall A.27b.nonempty \ A.27b \Rightarrow (\\ \forall V0f \in (A.27b^{A.27a}).(\forall V1P \in (2^{A.27b}).((p \ (ap \ (c.2EquantHeuristics.2EIS.REMOVABLE.QUANT.FUN \\ A.27a \ A.27b) \ V0f)) \Rightarrow ((\forall V2x \in A.27a.(p \ (ap \ V1P \ (ap \ V0f \ V2x)))) \Leftrightarrow \\ & (\forall V3x.27 \in A.27b.(p \ (ap \ V1P \ V3x.27))))) \end{aligned}$$